

Stability of Poisson Equilibria and Hamiltonian Relative Equilibria by Energy Methods

George W. Patrick

Mathematics and Statistics
University of Saskatchewan
Saskatoon, Saskatchewan, S7N 5E6
Canada
patrick@math.usask.ca

Mark Roberts

Mathematics and Statistics
University of Surrey
Guildford GU2 7XH
United Kingdom
m.roberts@surrey.ac.uk

Claudia Wulff*

Mathematics Institute
University of Warwick
Coventry CV4 7AL
United Kingdom
claudia@maths.warwick.ac.uk

January 2002

Abstract

We develop a general stability theory for equilibrium points of Poisson dynamical systems and relative equilibria of Hamiltonian systems with symmetries, including several generalisations of the Energy-Casimir and Energy-Momentum methods. Using a topological generalisation of Lyapunov's result that an extremal critical point of a conserved quantity is stable, we show that a Poisson equilibrium is stable if it is an isolated point in the intersection of a level set of a conserved function with a subset of the phase space that is related to the non-Hausdorff nature of the symplectic leaf space at that point. This criterion is applied to generalise the Energy-Momentum method to Hamiltonian systems which are invariant under non-compact symmetry groups for which the coadjoint orbit space is not Hausdorff. We also show that a G -stable relative equilibrium satisfies the stronger condition of being A -stable, where A is a specific group-theoretically defined subset of G which contains the momentum isotropy subgroup of the relative equilibrium.

*On leave from: Freie Universität Berlin

Contents

1	Introduction	2
2	Topology and Stability	5
3	Stability of Poisson Equilibria	6
3.1	Topological Tests	6
3.2	T_2 -Energy-Casimir Method	8
3.3	Examples	10
4	Stability of Hamiltonian Relative Equilibria	14
4.1	G -Stability by Topological Methods	15
4.2	Transverse Poisson Structures and Tame Generators	16
4.2.1	Coadjoint Actions	16
4.2.2	The Orbit Space P/G	19
4.3	T_2 -Energy-Momentum-Casimir Method	20
5	Beyond G-Stability	25
5.1	A -Stability	26
5.2	Split Momenta	27
5.3	Euclidean Invariant Hamiltonian Systems	28

1 Introduction

Energy methods for determining the stability of trajectories of Hamiltonian systems are based on the general principle that an equilibrium point is Lyapunov stable if it is a strict local minimum or maximum of a conserved function, such as the Hamiltonian itself. The *Energy-Casimir* and *Energy-Momentum methods* are extensions of this principle to, respectively, equilibrium points of Poisson systems and relative equilibria of Hamiltonian systems with symmetry. They originate with the work of Arnold on the stability of equilibria of incompressible fluids [2]. Since then they have been used very extensively in applications to rigid bodies [18, 20, 34], elasticity theory [21, 40, 43, 44], fluids [3, 4, 9, 14, 24], and vortex structures [16, 36, 39, 46]. In this paper we first present a topological generalisation of the energy method and then use this to obtain significant generalisations of the Energy-Casimir and Energy-Momentum methods.

The flow of a Poisson system on a Poisson manifold X generated by a Hamiltonian h preserves both h and the symplectic leaves of X . A point x_e is an equilibrium point of this flow if and only if it is a critical point of the restriction of h to the leaf $L(x_e)$ through x_e . If x_e is a local extremum of the restriction then the standard energy method implies that x_e is stable as an equilibrium point of the flow on $L(x_e)$. In this case we say that x_e is *leafwise stable*. In general x_e is not a critical point of h on the full space X . To test for stability on the whole of X the Energy-Casimir method supposes that there is a function C , the Casimir, which is constant on symplectic leaves and such that x_e is a critical point of $h + C$. Stability follows if this critical point is a local extremum.

When is it possible to find a Casimir C such that x_e is a critical point of $h + C$? One case is when x_e is a *regular* point of X , which means that locally the foliation into symplectic leaves is non-singular. Using this fact Arnold [3] and Libermann and Marle [22] show that if x_e is regular and is a local extremum of the restriction of h to the leaf $L(x_e)$, then x_e is stable for the full flow on X . Thus at regular points this test for leafwise stability is also a test for full stability. Examples show that this is not true in general, see [22, Exercise IV

15.10] and Examples 3.10, 3.11 and 4.20 of this paper. In such cases it is natural to ask whether there exists a space between $L(x_e)$ and X such that x_e is stable if it is an extremal point of the restriction of h to the intermediate space. In this paper we show that there is. More generally we answer a challenge posed by Weinstein [48] when, referring to the interaction between Poisson structures and stability, he wrote: “As yet there is no general theory for this kind of analysis”.

Most of the results in this paper are based on a topological generalisation of the energy method (Corollary 2.4) which generalises a lemma of Montaldi [31]. Corollary 2.4 is valid for a continuous flow on a locally compact topological space X which has conserved quantities with values in another topological space. In the case of Poisson systems the conserved quantities are the Hamiltonian h and the quotient map to the space of symplectic leaves. An equilibrium x_e is stable if the leafspace is Hausdorff at $L(x_e)$ and x_e is an isolated point in the fibre of the restriction of h to $L(x_e)$. Thus the condition that x_e be regular in the result of Arnold, Libermann and Marle can be relaxed to the leafspace being Hausdorff at $L(x_e)$. If the leafspace is not Hausdorff then h must isolate x_e in a larger subset $T_2(x_e)$ which depends only on the topology of the leafspace (Theorem 3.1).

We recover and generalise the Energy-Casimir method for Poisson equilibria: we will see that it suffices to make the assumptions of the Energy-Casimir method on a subset of the Poisson manifold X which contains $T_2(x_e)$. Example 3.13 shows that this improvement can succeed where the standard Energy-Casimir method fails. Moreover we identify a necessary condition for the Energy-Casimir method to apply, namely that the Poisson equilibrium must be *tame*. Our topological results provide stability tests which are even more general.

We now turn to the stability of relative equilibria. We consider a G -invariant Hamiltonian H on a symplectic manifold P with a momentum map $J : P \rightarrow \mathfrak{g}^*$ which commutes with the action of G on P and its coadjoint action on $\mathfrak{g}^*$, the dual of the Lie algebra $\mathfrak{g}$ of G . If the action of G on P is free and proper then the orbit space P/G is a Poisson manifold and criteria for the stability of Poisson equilibria can be lifted to criteria for the G -stability of relative equilibria. The symplectic leaves of P/G are just the Marsden-Weinstein reduced phase spaces and J induces a homeomorphism between the leafspace of P/G and the coadjoint orbit space $\mathfrak{g}^*/G$.

Leafwise stability of a relative equilibrium $p_e \in P$ means that it is stable to *momentum-preserving* perturbations of the initial condition, and is implied if the relative equilibrium is an extremal point of the reduced Hamiltonian. If the momentum $\mu_e = J(p_e)$ is regular, or more generally $\mathfrak{g}^*/G$ is Hausdorff at $G\mu_e$, then this condition also implies that p_e is G -stable (Remark 4.4). When the Hessian of the reduced Hamiltonian is definite and the momentum regular this was proved by Libermann and Marle [22], following earlier work of Arnold [3] and Marsden and Weinstein [28]. The result for $\mathfrak{g}^*/G$ Hausdorff is due to Montaldi [31].

The Energy-Momentum method is a convenient lifting of these criteria to the phase space. A point p_e is a relative equilibrium if and only if it is a critical point of the energy-momentum function $H_{\xi_e} = H - J_{\xi_e}$, where ξ_e is the *generator* of the relative equilibrium, an element of the Lie algebra of G , and $J_{\xi_e}(p) = \langle J(p), \xi_e \rangle$. The relative equilibrium is leafwise stable if the restriction of the Hessian $d^2 H_{\xi_e}(p_e)$ to a *symplectic normal space* N_1 is definite [27, 40, 42, 43, 44]. Full stability results were obtained by Patrick [35] for compact groups, and Ortega and Ratiu [33] and Lerman and Singer [19] under more general assumptions which still imply that $\mathfrak{g}^*/G$ is Hausdorff at $G\mu_e$. See also Corollary 4.13.

Some stability results for non-compact groups at momentum values μ_e where $\mathfrak{g}^*/G$ is not Hausdorff have been obtained by Leonard and Marsden [18] for semidirect products

of compact groups and vector spaces. They suggested that it is necessary to test for the definiteness of $d^2H_{\xi_e}(p_e)$ on a larger subspace of $T_{p_e}P$ than N_1 . In this paper we sharpen and generalise their results to arbitrary groups by lifting the general stability criteria for Poisson equilibria to conditions on H_{ξ_e} . In particular we identify subsets containing N_1 on which the definiteness of $d^2H_{\xi_e}(p_e)$ is sufficient to imply G -stability (Theorem 4.15 and Corollary 4.17).

The papers [19, 33, 35] all show, under various conditions which in particular imply that $\mathfrak{g}^*/G$ is Hausdorff at $G\mu$, that the definiteness of $d^2H_{\xi_e}(p_e)$ on N_1 implies that p_e is actually G_{μ_e} -stable, ie trajectories that start near p_e remain near $G_{\mu_e}p_e$, where G_{μ_e} is the coadjoint isotropy subgroup at μ_e . However numerical integration of an example by Leonard and Marsden [18] suggests that this is not always true, even for a regular momentum value μ_e . They prove that these relative equilibria are Γ -stable for a subgroup Γ which lies strictly between G_{μ_e} and G .

In this paper we introduce the general notion of A -stability for any subset of G and then show that any G -stable relative equilibrium is automatically A -stable where A is a ‘cone about G_{μ_e} ’ which can be made arbitrarily close to G_{μ_e} by restricting the perturbations from p_e to be sufficiently small (Theorem 5.2). In many cases this result can be improved by decomposing G_{μ_e} into the product of an (essentially) compact subgroup and a non-compact submanifold and showing that the ‘cone’ only needs to be taken about the non-compact part. (Theorem 5.3). As a corollary we obtain a generalisation of the results of [19, 33, 35] on G_{μ_e} -stability (Corollary 5.4). Unlike the previous results we do not require a Hessian condition to be satisfied, only that the relative equilibrium is G -stable.

Taken together our results on the stability of Poisson equilibria and the G and A -stability of relative equilibria provide generalisations of all previous results for finite dimensional Poisson and Hamiltonian systems that we are aware of. We believe that our most primitive topological tests for stability (Theorems 3.1 and 4.2) are the sharpest possible general results, and that they fully explain the interaction between stability properties and Poisson structures. In this paper we restrict attention to free Hamiltonian group actions for simplicity only; the extensions to general proper actions are contained in [37].

We end this introduction with a guide to the paper. Section 2 contains the fundamental topological ideas that underly the rest of the paper. Topological and derivative tests for the stability of Poisson equilibria are described in Sections 3.1 and 3.2, respectively. Section 3.2 introduces the machinery of *smoothings* that is used to make sense of derivatives of functions on singular sets and the related notion of *tame generators*. It also describes the role Casimirs play in stability theory and their limitations. The topological and derivative tests for the G -stability of Hamiltonian relative equilibria are contained in Sections 4.1 and 4.3, respectively. The intermediate Section 4.2 introduces *transverse Poisson structures* as tools for describing local leafspace topology and uses this to discuss some special cases. Towards the end of Section 4.3 we discuss how the stability criteria apply to relative equilibria of systems that are invariant under actions of the Euclidean groups $SE(2)$ and $SE(3)$ and how our theory ‘explains’ an example of Libermann and Marle [22]. Section 5 is devoted to A -stability: Section 5.1 gives the general result, Section 5.2 the improvements obtained using Lie group decompositions and Section 5.3 the application to Euclidean groups.

2 Topology and Stability

Let X and Y be a topological spaces and $f: X \rightarrow Y$ a continuous map. In this section we give a stability criterion for equilibria of continuous flows on X which preserve the fibres of f . This is a corollary of a ‘confinement’ result for general continuous curves preserving f . The confinement result is inspired by Lemma 1.4 of Montaldi [31] in which Y is Hausdorff.

We begin with a measure of the extent to which a space is not Hausdorff at a point.

Definition 2.1 Let Y be a topological space and $y \in Y$. Define

$$T_2(y) \equiv \{ y' \in Y : U \cap U' \neq \emptyset \text{ for all neighbourhoods } y \in U \subseteq Y \text{ and } y' \in U' \subseteq Y \}.$$

We say that a topological space X is *Hausdorff* at $x \in X$ if $T_2(x) = \{x\}$. The following is frequently useful and easily proved.

Proposition 2.2

1. Let X and Y be topological spaces and $(x, y) \in X \times Y$. Then $T_2(x, y) = T_2(x) \times T_2(y)$.
2. If X , Y_1 and Y_2 are topological spaces and $f_i: X \rightarrow Y_i$, $i = 1, 2$, are continuous, then $(f_1 \times f_2)^{-1}(T_2(y_1, y_2)) = f_1^{-1}(T_2(y_1)) \cap f_2^{-1}(T_2(y_2))$. In particular, if Y_2 is Hausdorff then $(f_1 \times f_2)^{-1}(T_2(y_1, y_2))$ is the y_2 level set of $f_2|_{f_1^{-1}(T_2(y_1))}$.

The next result gives a sufficient condition for continuous curves in X preserving f and starting near a point x to be confined to small neighbourhoods of x .

Lemma 2.3 (Topological Stability Lemma) Let X and Y be topological spaces, $f: X \rightarrow Y$ a continuous map, $x \in X$ and $y = f(x)$. Assume that:

1. X is locally compact at x ;
2. There exists a neighbourhood U of x in X such that $f^{-1}(T_2(y)) \cap U = \{x\}$.

Then for every neighbourhood U of x there is a neighbourhood V of x such that if $c: [0, 1] \rightarrow X$ is a continuous curve for which $f \circ c$ is constant and $c(0) \in V$, then $c(t) \in U$ for all $t \in [0, 1]$.

If the second condition holds we say that x is an *isolated point* of $f^{-1}(T_2(y))$.

Proof. We will prove the lemma by showing that if f -constant curves are not confined to small neighbourhoods of x then x is not an isolated point in $f^{-1}(T_2(y))$. Let $\mathcal{B}$ be a neighbourhood base at x consisting of compact sets. For any $U \in \mathcal{B}$ and any $V \in \mathcal{B}$ such that $V \subseteq U$, let $c_{U,V}: [0, 1] \rightarrow X$ be a continuous curve such that $f \circ c$ is constant, $c_{U,V}(0) \in V$, and $c_{U,V}(1) \notin U$. By connectedness of $[0, 1]$ there exists $t_{U,V} \in (0, 1]$ such that $x_{U,V} = c_{U,V}(t_{U,V}) \in \partial U$.

For fixed $U \in \mathcal{B}$ the set $\{x_{U,V}\}_{V \in \mathcal{B}}$ is a net in ∂U by reverse inclusion of the V ’s. Since ∂U is compact there is a subnet, $\{x_{U,V_\lambda}\}$, which converges to a point $z_U \in \partial U$. The continuity of f implies that $\{f(x_{U,V_\lambda})\}$ converges to $f(z_U)$. Since the curves $c_{U,V}$ preserve f we also have:

$$f(x_{U,V_\lambda}) = f(c_{U,V_\lambda}(t_{U,V_\lambda})) = f(c_{U,V_\lambda}(0)).$$

The net $c_{U,V_\lambda}(0)$ converges to x and so $f(x_{U,V_\lambda})$ also converges to $y = f(x)$. Thus, every neighbourhood of $f(z_U)$ meets every neighbourhood of y , and so $f(z_U) \in T_2(y)$. This is the required contradiction, since $z_U \in f^{-1}(T_2(y))$ and $\{z_U\}_{U \in \mathcal{B}}$ is a net converging to x . ■

Now consider a continuous flow $\phi: X \times \mathbb{R} \rightarrow X$ on X which preserves the fibres of f , $f(\phi_t(x)) = f(x)$, and which has an equilibrium point at $x \in X$. The equilibrium point is *stable* if for every open neighbourhood U of x in X there exists a neighbourhood V of x such that if $x \in V$ then $\phi_t(x) \in U$ for all t .

Corollary 2.4 *Let X and Y be topological spaces, X locally compact, and $f: X \rightarrow Y$ a continuous map. Let x be an equilibrium point of a continuous flow on X which preserves f and $y = f(x)$. Then x is stable if it is an isolated point in $f^{-1}(T_2(y))$.*

The examples in Section 3.3 show that Corollary 2.4 is false in general if x is only isolated in $f^{-1}(y)$. The following example shows that it is necessary to assume that X is locally compact:

Example 2.5 For $x = (q_n, p_n)_{n \geq 0} \in X = \ell^2(\mathbb{R}) \times \ell^2(\mathbb{R})$ let

$$h(x) = \frac{1}{2} \sum_{n=0}^{\infty} p_n^2 + \frac{1}{2} \sum_{n=0}^{\infty} 4^{-n} q_n^2.$$

Then h is differentiable on X and $h(0) = 0$. Clearly $T_2(0) = \{0\}$ and 0 is isolated in $h^{-1}(0) \subset X$. The standard linear symplectic structure $J((q_n, p_n)) = (p_n, -q_n)$ gives the Hamiltonian system $\dot{x} = Jdh(x)$ and defines a flow ϕ_t on X . For $m \geq 1$ the solution of this Hamiltonian system with initial value such that $q_m = 2^{-m}$, $q_n = 0$ for $n \neq m$ and $p_n = 0$ for all n , has $p_m = -1$ at time $t = 2^{m-1}\pi$, as is easily verified. Hence 0 is an unstable equilibrium.

The requirement that X is locally compact implies that Corollary 2.4 can not be applied directly to deduce the nonlinear stability of equilibria of partial differential equations. Indeed, it is well-known that for partial differential equations positivity of the second variation need not imply stability of an equilibrium, a phenomenon which is related to the non-equivalence of norms on Banach spaces [5].

3 Stability of Poisson Equilibria

In this section we apply the topological stability result of Section 2 to equilibria of Poisson systems to obtain generalisations of the Energy-Casimir method.

3.1 Topological Tests

A finite dimensional Poisson manifold X is partitioned into immersed submanifolds by its symplectic leaves, which without loss of generality we will always assume to be connected. Define two points in X to be equivalent if they belong to the same symplectic leaf, let Z be the quotient of X by this equivalence relation and $L: X \rightarrow Z$ the quotient map. We will regard the symplectic leaf $L(x)$ through x as both a subset of X and as a point in Z .

A function $h : X \rightarrow \mathbb{R}$ generates a vector field on X which is uniquely defined by the requirement that $\dot{f} = \{h, f\}$ for all differentiable functions f on X . The flow ϕ_t of X preserves the fibres of both h and L , and hence those of the product map $f = L \times h : X \rightarrow Y = Z \times \mathbb{R}$. We can therefore apply the topological stability result Corollary 2.4.

First note that $x_e \in X$ is an equilibrium point of ϕ_t if and only if it is an equilibrium point of the restriction of the flow to the invariant submanifold $L(x_e)$ or, equivalently, if and only if the restriction of h to $L(x_e)$ has a critical point at x_e . An equilibrium x_e is said to be *leafwise stable* if it is stable for the restricted flow on $L(x_e)$. Clearly stability in the full space X implies leafwise stability.

For $x \in X$, let $T_2(x) = L^{-1}(T_2(L(x)))$. Note that $L(x) \subseteq T_2(x)$. More generally, for any open neighbourhood U of x in X let $L^U(x)$ denote the symplectic leaf of U through x . This is the connected component of $L(x) \cap U$ which contains x . Denote the space of these symplectic leaves by Z^U and the corresponding quotient map by $L^U : U \rightarrow Z^U$. Define

$$T_2^U(x) = (L^U)^{-1}(T_2(L^U(x))),$$

where $T_2(L^U(x))$ is taken in Z^U . The set $T_2^U(x)$ is contained in $T_2(x) \cap U$, but may be strictly smaller, for example if $L(x)$ accumulates on itself at x .

Recall that a point x_e in a Poisson manifold X is said to be *regular* (or *minimal*) if there exists an open neighbourhood U of x_e in X such that $\dim L(x) = \dim L(x_e)$ for all x in U . The set of regular points is open and dense in X since in local coordinates it corresponds to the set where the matrix of the Poisson tensor has locally constant rank.

The following result provides topological conditions for leafwise stability and stability and states that for regular equilibria these conditions coincide.

Theorem 3.1 *Let x_e be an equilibrium point of the flow generated by a Hamiltonian h on a Poisson manifold X , and U be an open neighbourhood of x_e . Then x_e is*

1. *leafwise stable if there is an open neighbourhood U of x_e in X such that $h^{-1}(h(x_e)) \cap L^U(x_e) = \{x_e\}$; and*
2. *stable if there is an open neighbourhood U of x_e in X such that $h^{-1}(h(x_e)) \cap T_2^U(x_e) = \{x_e\}$.*

Moreover, if x_e is a regular point of X then there is an open neighbourhood U of x_e in X such that $T_2^U(x_e) = L^U(x_e)$ and part 1 implies stability.

These statements remain true if h is replaced by any conserved quantity with values in a Hausdorff space.

Proof.

1. This follows from Corollary 2.4 with X replaced by $L^U(x_e)$ and f by h . Clearly, $T_2(h(x_e)) = \{h(x_e)\}$ and the hypothesis says that x_e is an isolated point in the $h(x_e)$ level set of the restriction of h to $L^U(x_e)$.
2. Apply Corollary 2.4 to the neighbourhood U and map $f = L^U \times h$, noting that by Proposition 2.2 we have $T_2(y) = T_2(L^U(x_e), h(x_e)) = T_2(L^U(x_e)) \times \{h(x_e)\}$ and so $f^{-1}(T_2(y)) = h^{-1}(h(x_e)) \cap T_2^U(x_e)$.

If x_e is regular then there exists an open neighbourhood U of x_e in X for which the symplectic leaves provide a regular foliation [47, Corollary 2.3]. The quotient of U by this foliation is Hausdorff and so $T_2^U(x_e)$ is equal to $L^U(x_e)$. ■

In Example 3.10 we show that there is an open subset of Hamiltonians on $\mathfrak{sl}(2)^* \cong \mathbb{R}^3$ for which Theorem 3.1 implies that the origin is a stable equilibrium, and another open set for which the origin is leafwise stable but not stable.

The next result is a simple corollary of Theorem 3.1 for a very special case.

Corollary 3.2 *Let x_e be an equilibrium point of a Hamiltonian h on the Poisson manifold X . Suppose that $T_2^U(x_e)$ is a one-dimensional submanifold of X . Then x_e is stable if $dh(x_e)$ is nonzero on $T_{x_e}(T_2^U(x_e))$.*

Proof. If $dh(x_e)$ is nonzero on $T_{x_e}(T_2^U(x_e))$ then the level set of h containing x_e intersects $T_2(x_e)$ transversely and hence in an isolated point. ■

See Example 3.9 for an illustration of this result. The hypotheses of Corollary 3.2 can only be satisfied if x_e is a zero-dimensional symplectic leaf, in which case x_e is trivially a leafwise stable equilibrium for any Hamiltonian. In Example 3.11 we show that the assumption that $T_2^U(x_e)$ be one-dimensional cannot be replaced by the assumption that $L(x_e)$ has codimension one in $T_2^U(x_e)$. Results analogous to Corollary 3.2, in the sense that $dh(x_e) \neq 0$ and stability can be determined through a condition on $dh(x_e)$, can also be obtained for some cases in which $T_2(0)$ is higher dimensional but singular at x_e . See Example 3.10.

3.2 T_2 -Energy-Casimir Method

We now discuss conditions on the derivative and Hessian of h at x_e for stability to hold. If x_e is an equilibrium, and so the restriction of h to $L(x_e)$ has a critical point at x_e , then it is leafwise stable if the second derivative of the restriction is definite. The following result, a special case of Theorem 3.1, states that for generic points in X this condition also implies that x_e is stable.

Proposition 3.3 [22, Theorem III.12.4] *If an equilibrium x_e is regular and the second derivative of the restriction of the Hamiltonian h to $L(x_e)$ is positive or negative definite, then x_e is both leafwise stable and stable.*

If $T_2^U(x_e)$ contains a two dimensional manifold passing through x_e on which $dh(x_e) \neq 0$ then the level set of h in $T_2^U(x_e)$ through x_e contains a smooth curve and so h does not isolate x_e in $T_2^U(x_e)$. In such a case stability cannot be concluded from the mechanism of confinement by energy level sets. However, if $T_2^U(x_e)$ is a manifold, if the first derivative of h vanishes on $T_2^U(x_e)$ and if the second derivative of the restriction of h to $T_2^U(x_e)$ is definite, then h again isolates x_e and x_e is stable. We call Poisson equilibria for which the first derivative of h satisfies this condition, and which are therefore amenable to Hessian type stability tests, *tame*. To extend this to cases for which $T_2^U(x_e)$ is not a manifold, we first define the notions of *tangent space* and *smoothing* of a singular set.

Definition 3.4 Let M be a manifold, $S \subseteq M$ and $m \in S$.

1. The *tangent space* $T_m S$ of S at m is the subset of $T_m M$ consisting of the derivatives $c'(0)$ of all C^1 curves $c(t)$ in M with $c(0) = m$ and $c(t) \in S$ for $t \geq 0$.
2. A *smoothing* of S at m is a finite number of submanifolds $B_i \subseteq M$ such that $S \subseteq \bigcup_{i=1}^n B_i$ and $T_m B_i \subseteq \text{span}(T_m S)$ for $1 \leq i \leq n$.

If $T_2^U(x_e)$ is a manifold then clearly it is its own smoothing. Examples 3.10 and 3.12 feature some smoothings of singular $T_2^U(x_e)$ sets. *Weak smoothings* which satisfy the first condition of 2., but not the second, always exist since it is possible to take $n = 1$ and $B_1 = M$. In Example 3.15 the T_2 -set does not have a smoothing.

We now define tame equilibria at points with arbitrary T_2 -sets.

Definition 3.5 Let X be a Poisson manifold.

1. A *generator* at x is an element $\xi \in T_x^*X$ which annihilates $T_x L(x)$.
2. A generator ξ at x is *tame* if it annihilates $T_x(T_2^U(x))$. Generators that are not tame are said to be *wild*.
3. The generator of an equilibrium x_e of a Poisson system with Hamiltonian h is $dh(x_e)$.
4. An equilibrium is tame if its generator is tame, and wild otherwise.

We will notationally suppress reference to U even though the property of being tame is U dependent. The set of tame generators at x is a vector subspace of T_x^*X . If $\xi \in T_x^*X$ is tame then it annihilates every smoothing $\{B_i\}_{i=1}^n$ of $T_2(x)$, ie ξ annihilates $T_x B_i$ for all i .

If x is regular then $T_2^U(x) = L^U(x)$ for some neighbourhood U and every generator is tame. In Corollary 3.2 the hypothesis that $dh(x_e)$ is nonzero on $T_{x_e}(T_2^U(x_e))$ implies that $dh(x_e)$ is wild. In Example 3.10, for which $T_2(0)$ is a cone, $T_0(T_2(0))$ spans the whole of $X = \mathbb{R}^3$ and so every nonzero derivative at 0 is wild.

A *Casimir* on U is a continuous function $C : U \rightarrow \mathbb{R}$ which is constant on the symplectic leaves of U . This condition implies that C is also constant on every set $T_2^U(x)$ for every $x \in U$. Casimirs are conserved quantities along integral curves contained in U , since any Poisson flow preserves the symplectic leaves. Since Casimirs are constant on $T_2^U(x)$, the derivatives of smooth Casimirs at x are tame. It follows that if x_e is an equilibrium and C is a smooth Casimir at x_e , then $dh(x_e)$ is tame if and only if $d(h + C)(x_e)$ is tame.

Theorem 3.6 (Poisson T_2 -Energy-Casimir Method) *Let x_e be a tame equilibrium point of the Hamiltonian h . Let $\{B_i\}_{i=1}^n$ be a smoothing of $T_2^U(x_e)$ at x_e . Then x_e is stable if for each i there is a smooth Casimir C_i such that the Hessian $d^2((h + C_i)|_{B_i})(x_e)$ is positive or negative definite on $T_{x_e}B_i$.*

Proof. Set $\hat{h} = (h + C_1, \dots, h + C_n)$. Since $T_2^U(x_e) \subseteq \bigcup_{i=1}^n B_i$, by Theorem 3.1 it suffices to show that $\hat{h}$ isolates x_e on $\bigcup_{i=1}^n B_i$. As there are only finitely many B_i , this follows if $h + C_i$ isolates x_e on each B_i . This in turn is implied by the definiteness of the Hessians and the Morse lemma. ■

The standard Energy-Casimir method, see eg [26], states that if x_e is an equilibrium point of the flow generated by a Hamiltonian h and there exists a smooth Casimir C such that x_e is a critical point of $h + C$ on the whole of X and $d^2(h + C)(x_e)$ is definite, then x_e is stable. Theorem 3.6 is a strict generalisation of this because it only requires x_e to be a definite critical point for a function on a subset of X . Example 3.13 describes a Poisson system for which the theorem can be used to prove stability, though the standard Energy-Casimir method fails.

The standard Energy-Casimir method and its generalisation, Theorem 3.6, can only be applied to tame equilibria, but the topological stability result Theorem 3.1 can sometimes

be applied to wild equilibria. Examples include the cases covered by Corollary 3.2 and Examples 3.9 and 3.10.

Smoothings and Casimirs are both implements designed to handle singularities of $T_2^U(x_e)$ for the purpose of constructing Hessian tests for stability. Fine smoothings can eliminate the necessity of including Casimirs when calculating the Hessians, while coarse smoothings may require their inclusion. Example 3.12 illustrates the play available in choosing the smoothings versus Casimirs.

If there exists a smoothing such that $T_2^U(x_e) = \bigcup_{i=1}^n B_i$ (for example, if $T_2^U(x_e)$ is a submanifold) then every Casimir is constant on each of the B_i . In this case the second derivatives of the restrictions of the Casimirs to the B_i vanish, and the inclusion of the Casimirs in the Hessians in Theorem 3.6 is unnecessary. However, Example 3.10 shows that inclusion of Casimirs can be necessary in cases when $T_2^U(x_e)$ is singular.

More generally, the inclusion of the Casimir C_i is unnecessary when all Casimirs restricted to B_i have vanishing second derivative at x_e . This follows if $B_i \subseteq T_2^U(x_e)$ but can also be implied by an infinitesimal relation between the smoothing and the set $T_{x_e}^U(x_e)$, as follows. For any Casimir C we have $d^2C(x_e)(v, v) = 0$ for all $v \in T_{x_e}(T_2^U(x_e))$, since Casimirs are constant on curves in $T_2^U(x_e)$. Regarding the symmetric bilinear form $d^2C(x_e)$ as a linear map on the tensor product $T_{x_e}B_i \otimes T_{x_e}B_i$, it follows that $d^2C(x_e)$ vanishes on the whole of $T_{x_e}B_i$ if

$$\text{span} \{ v \otimes v : v \in T_{x_e}(T_2^U(x_e)) \cap T_{x_e}B_i \} = T_{x_e}B_i \otimes T_{x_e}B_i. \quad (3.1)$$

Consequently, in Theorem 3.6, the inclusion of C_i is unnecessary for any i such that this spanning condition holds. An application of this is given in Example 3.13.

Remark 3.7 If $\{B_i\}_{i=1}^n$ is only a weak smoothing of $T_2(x_e)$, but there exist Casimirs C_i such that for $i = 1, \dots, n$ we have $d(h + C_i)(x_e)|_{B_i} = 0$ and the Hessians $d^2(h + C_i)(x_e)|_{B_i}$ are definite, then x_e is again stable. This can be deduced as in the proof of Theorem 3.6. The standard Energy-Casimir method is recovered by taking $n = 1$ and $B_1 = M$.

3.3 Examples

Here we collect together a number of examples to illustrate the stability theory described above. Many of the examples are of equilibria on Poisson spaces which are duals of Lie algebras, for which the symplectic leaves are coadjoint orbits. Other examples are Poisson structures on $X = \mathbb{R}^3$ with Poisson brackets of the form:

$$\{h, f\} = \nabla A \cdot (\nabla h \times \nabla f) \quad (3.2)$$

where $A = A(x, y, z)$ is a smooth function. The vector field generated by a Hamiltonian h is

$$\dot{x} = \nabla A \times \nabla h.$$

We may assume that $A(0) = 0$. For these structures A is a Casimir since $\{A, f\} = 0$ for all functions h , and the two dimensional symplectic leaves of X are the connected components of the level sets of regular values of A . Each critical point x of A is also a symplectic leaf since there $\{h, f\}(x) = 0$ for all smooth functions h and f . For any open neighbourhood U of 0 in X the set $T_2^U(x)$ is contained in the connected component of $A^{-1}(0) \cap U$ which contains 0. If $A = \frac{1}{2}(x^2 + y^2 + \epsilon z^2)$ then X is isomorphic to the dual of the Lie algebra, $\mathfrak{g}^*$, of

$SO(3)$ ($\epsilon = 1$), $SE(2)$ ($\epsilon = 0$) or $SL(2, \mathbb{R})$ ($\epsilon = -1$) with their standard Poisson structures. These three Poisson structures are described in [26] and are also used in [48] to illustrate the interdependence of stability and Poisson structure.

Example 3.8 If $A = \frac{1}{2}(x^2 + y^2 + z^2)$, and so $X \cong so(3)^*$, then for any neighbourhood U of 0 we have $L(0) = T_2^U(0) = \{0\}$ and so 0 is an isolated point of $h^{-1}(h(0)) \cap T_2^U(0)$, and hence a stable equilibrium, for any Hamiltonian h . Every generator at 0 is tame. Every point $x_e \neq 0$ is regular, since the symplectic leaf through each nearby point (a sphere) is two dimensional. Thus all generators at these points are also tame. In fact every generator is tame at every point in $\mathfrak{g}^*$ for any compact group, since in this case the symplectic leaves are the coadjoint orbits of G on $\mathfrak{g}^*$ and the quotient space $\mathfrak{g}^*/G$ is always Hausdorff.

Example 3.9 If $A = \frac{1}{2}(x^2 + y^2)$ then $X \cong se(2)^*$. The symplectic leaves are points on the z -axis and cylinders of nonzero radius about the z -axis.

If x_e is a point on the z -axis then $T_2^U(x_e)$ is the intersection of the z -axis with U , a one dimensional manifold. The set of tame generators at x_e is equal to the subspace $\mathbb{R}^2 \subseteq se(2)$ consisting of infinitesimal translations. If $dh(x_e)$ is tame the point x_e is stable if the restriction of h to the z -axis is positive or negative definite. Generators containing nonzero rotational components are wild. However in this case this implies that $dh(x_e)$ is nonzero on $T_{x_e}(T_2^U(x_e)) = \{x = y = 0\}$ and so 0 is again stable by Corollary 3.2.

If x_e does not lie on the z -axis then it is regular and every generator is tame.

Example 3.10 Next consider $A = \frac{1}{2}(x^2 + y^2 - z^2)$, giving $X \cong sl(2, \mathbb{R})^*$. The symplectic leaves are the connected components of the hyperboloids $A = a$ for $a \neq 0$, the connected components on the complement of 0 in the cone $A = 0$, and the origin itself. Every nonzero point is regular and all generators at these points are tame.

At the origin, $T_2^U(0)$ is the intersection of the cone $A = 0$ with U . Since $T_0(T_2^U(0))$ spans the whole of $\mathbb{R}^3$ the only tame generator is $\xi = 0$. Nevertheless Hamiltonians with wild generators $dh(0) \neq 0$ can again have stable equilibria at 0. Theorem 3.1 implies that 0 is stable if $dh(0)$ ‘points into the cone $A = 0$ ’, so that $\text{ann}(dh(0))$ intersects the cone only at the origin. If $dh(0)$ points out of the cone then the intersection is (infinitesimally) a pair of lines and the equilibrium is unstable. The instability follows from the fact that if $dh(0) = (\xi_1, \xi_2, \xi_3)$ the linearised equations of motion at the origin have eigenvalues 0 and $\pm\sqrt{\xi_1^2 + \xi_2^2 - \xi_3^2}$. Since 0 is trivially leafwise stable for any Hamiltonian this is an example of an equilibrium that is leafwise stable but not stable.

Now consider the tame case $dh(0) = 0$. For a smoothing we have to take $B = U$, a full neighbourhood of the origin. Without including Casimirs we can only use Theorem 3.6 to conclude stability if $d^2h(0)$ is definite. However by using the Casimir A it can be seen that, for example, the Hamiltonian $ax^2 + by^2 + cz^2$ has a stable equilibrium whenever $c > -a$ and $c > -b$. Thus in general Casimirs can not be dispensed with completely.

Example 3.11 Let X denote the Poisson space $se(2)^* \times \mathbb{R}^2$, with coordinates x, y, z on $se(2)^*$, as in Example 3.9, and coordinates p, q on $\mathbb{R}^2$ satisfying $\{p, q\} = 0$. Then the symplectic leaf through 0 is the two dimensional manifold $L(0) = \{x = y = z = 0\}$ and $T_2^U(0)$ is the intersection of U with the three dimensional manifold $\{x = y = 0\}$. The set of

generators can be identified with $T_0^*(se(2)^*) \cong se(2)$ and the tame generators are the pure translations, as in Example 3.9.

However in contrast to Example 3.9, Corollary 3.2 does not apply, and 0 can be unstable when $dh(0)$ is wild. To see this consider the Hamiltonian $h = az - qy + \frac{1}{2}(q^2 + p^2)$. This has a wild generator at 0. Its restriction to the symplectic leaf has a strict local minimum at 0, so 0 is leafwise stable. The equations of motion are:

$$\dot{x} = ay, \quad \dot{y} = -ax, \quad \dot{z} = -qx, \quad \dot{q} = p, \quad \dot{p} = -q + y.$$

When $a = 1$ for arbitrarily small δ the curve

$$x = \delta \sin t, \quad y = \delta \cos t, \quad z = \frac{\delta^2}{8}(t \sin 2t - \sin^2 t - t^2), \quad q = \frac{\delta}{2}t \sin t, \quad p = \frac{\delta}{2}(\sin t + t \cos t)$$

is a solution which leaves any neighbourhood of 0 for t sufficiently large. This instability is already present in the linearised equations at 0 and is caused by the 1 : 1 resonance between the x, y and q, p frequencies.

Example 3.12 Let $X = \mathbb{R}^3$ with the Poisson structure given by (3.2) with $A = \frac{1}{2}(x^2 - y^2)$. The set $T_2^U(0)$ is the intersection of U with the union of the two planes $x = \pm y$. The only tame generator is $\xi = 0$. One smoothing of $T_2^U(0)$ is provided by the two planes themselves: $B_1 = \{x = y\}$, $B_2 = \{x = -y\}$. Another is obtained by taking a whole open neighbourhood U of 0.

Consider the Hamiltonian $h = ax^2 - by^2 + z^2$. The restriction of h to each B_i is $(a-b)x^2 + z^2$ and so 0 is stable by Theorem 3.6 if $a > b$, without using any Casimirs. Alternatively the Casimir A can be used on the whole of U :

$$h - 2\lambda A = (a - \lambda)x^2 - (b - \lambda)y^2 + z^2$$

and so if $a > b$ then taking $a > \lambda > b$ gives stability by the Energy-Casimir method.

Example 3.13 This is an example of an equilibrium which is stable by Theorem 3.6, but for which the standard Energy-Casimir method fails. Again let $X = \mathbb{R}^3$ with the Poisson structure given by (3.2), but with $A = (a^2x^2 - y^2)y$ where $a \neq 0$. The set $T_2^U(0)$ is the intersection of U with the union of the three planes $y = 0$ and $y = \pm ax$. The only tame generator is $\xi = 0$. The restriction of the Hamiltonian $h = x^2 - y^2 + z^2$ to $y = 0$ isolates 0 in $y = 0$, while its restrictions to $y = \pm ax$ isolate 0 if $|a| \leq 1$. Thus 0 is stable by Theorem 3.6 if $|a| \leq 1$. However any Casimir on X must satisfy $d^2C(0) = 0$ since, as v varies over the three planes forming the set $T_2(0)$, the vectors $v \otimes v$ span the symmetric part of $\mathbb{R}^3 \otimes \mathbb{R}^3$. Thus the standard Energy-Casimir method can not be used to deduce stability.

Example 3.14 This example shows that the use of $T_2^U(x_e)$ instead of the larger set $T_2(x_e)$ is necessary when the symplectic leaves accumulate upon themselves. Let $\hat{X} = \mathbb{R}^3$ with the Poisson bracket (3.2) with $A = x - ay$. Let

$$\hat{h} = \frac{1}{2}(z^2 + (\sin y - a \sin x)^2 - (\sin x - a \sin y)^2),$$

and let $\hat{x}_e = 0$. The action of $\mathbb{Z} \times \mathbb{Z}$ on $\hat{X}$ by $(b_1, b_2) \cdot (x, y, z) = (x + 2\pi b_1, y + 2\pi b_2, z)$ is Poisson and $\hat{h}$ is invariant. Let $X \equiv \hat{X}/\mathbb{Z} \times \mathbb{Z}$ be the Poisson quotient, and h and x_e be the projections of $\hat{h}$ and $\hat{x}_e$ to the quotient. Let a be irrational. Then the symplectic leaf through x_e is the projection of the plane $x = ay$ in $\hat{X}$. This is dense in X since it is the product of a densely winding line on a 2-torus and $\mathbb{R}$. So $T_2(x_e) = X$ and Theorem 3.6 fails to show that x_e is stable, since the Hessian of h is indefinite on X . However, by taking U to be the projection of $(-r, r) \times (-ar, ar) \times \mathbb{R}$ for sufficiently small r , the set $T_2^U(x_e)$ becomes the product of the projection of the *line segment* $y = ax$, $x \in (-r, r)$, and $\mathbb{R}$, which is not dense. The Hessian of h restricted to $T_2^U(x_e)$ is definite since $\hat{h} = \frac{1}{2}(z^2 + (1 - a^2)^2 y^2) + h.o.t.$ on the plane $x = ay$ in $\hat{X}$.

Example 3.15 Let $X = sl(3; \mathbb{R})^*$. Using the Killing form we can identify X with $sl(3; \mathbb{R})$, and hence with the space of traceless 3×3 real matrices. Let x_e denote the *subregular* nilpotent matrix with $(x_e)_{12} = 1$ and all other entries equal to zero. We claim that $T_2^U(x_e)$ does not have a smoothing for any neighbourhood U . The coadjoint orbit $L(x_e)$ has codimension four. Let Σ be a four-dimensional section through x_e transverse to $L(x_e)$. This has an induced Poisson structure that is described in Section 4.2 below and $T_2^U(x_e)$ is isomorphic to the product of a neighbourhood of x_e in $L(x_e)$ and $T_2^{U_\Sigma}(x_e)$ where U_Σ is a neighbourhood of x_e in Σ . We will show that $T_2^{U_\Sigma}(x_e)$ does not have a smoothing for any neighbourhood U_Σ .

Let $\pi : X \rightarrow \mathbb{R}^2$ denote the mapping defined by generators of the ring of invariants of the coadjoint action on X , ie the symmetric polynomials of the eigenvalues of the matrices. Let π_Σ be the restriction of π to Σ . Since the generators are Casimirs the set $T_2^{U_\Sigma}(x_e)$ is contained in the two-dimensional fibre of π_Σ through x_e . Brieskorn has shown that this fibre has a simple singularity of type A_2 [8, 45], which means that it is diffeomorphic to the variety defined by $x^2 + y^2 + z^3 = 0$. The tangent space to this variety at 0 (in the sense of Definition 3.4) is just the non-negative z -axis, ie $T_{x_e}(T_2^{U_\Sigma}(x_e)) = \{(0, 0, z) : z \geq 0\}$. However the fibre $\pi_\Sigma^{-1}(\pi_\Sigma(x_e))$ contains the intersection of Σ with the coadjoint orbit through the *regular* nilpotent matrix defined by $(x_e)_{12} = (x_e)_{23} = 1$ and all other entries equal to zero. This intersection is a symplectic leaf of Σ of dimension two which, by Jordan normal form theory, contains x_e in its closure. It must therefore be contained in $T_2^{U_\Sigma}(x_e)$. Thus a smoothing $\{B_i\}_{i=1}^n$ must have one B_i at least of dimension 2 and it is not possible that $T_{x_e} B_i \subset \text{span } T_{x_e}(T_2^{U_\Sigma}(x_e))$ for each i .

Example 3.16 This example shows that the assumption in Theorem 3.6 that the equilibrium is tame is essential, even for Poisson systems that are reductions of Hamiltonian systems of physically recognisable forms. Take $P = T^*(SE(2)^n)$ and $G = SE(2)^n$ acting by the cotangent lift of its left action on itself. Then $X = P/G$ is the Lie algebra dual $(se(2)^*)^n \cong (\mathbb{R}^3)^n$, and the generic coadjoint orbits are products of cylinders. In cylindrical coordinates (r_i, θ^i, z_i) the Poisson bracket is

$$\{f, g\} = \sum_{i=1}^n \left(\frac{\partial f}{\partial z_i} \frac{\partial g}{\partial \theta^i} - \frac{\partial g}{\partial z_i} \frac{\partial f}{\partial \theta^i} \right).$$

Consider Hamiltonians $h : P/G \rightarrow \mathbb{R}$ of the form

$$h = \sum_{i=1}^n \frac{1}{2} z_i^2 + r_i^2 F_i(\theta, z) \quad (3.3)$$

for smooth functions F_i on the product of n cylinders, $z = (z_1, \dots, z_n)$, $\theta = (\theta^1, \dots, \theta^n)$. Then the corresponding Poisson system is given by

$$\dot{r}_i = 0, \quad \dot{z}_i = -\partial_{\theta^i} h, \quad \dot{\theta}^i = \partial_{z_i} h, \quad i = 1, \dots, n.$$

For $i = 1, \dots, n$ the coordinate functions r_i are Casimirs that parametrise the Hamiltonian (3.3). Any point with all $r_i = 0$ is a symplectic leaf, and hence a (leafwise stable) equilibrium. The full stability of these equilibria corresponds to stability under small perturbation to nonzero r_i . These are perturbations of a completely integrable system, and so when $n \geq 3$ Arnold drift in the variables z_i is expected [1]. For $n = 3$ Bessi [7] has constructed functions F_i for which instability occurs near $z_1 = z_3 = 0$ and $z_2 = 2$. The T_2 -set corresponding to any equilibrium with all $r_i = 0$ is the set $\{(r_i, \theta^i, z_i) : r_1 = \dots = r_n = 0\}$ and dh restricted to the T_2 -set is not zero unless all z_i are zero. Thus the equilibrium is wild for $z \neq 0$.

We note that for $n = 1$ stability follows from Corollary 3.2, and that for $n = 2$ we essentially have a family of two-degree of freedom systems (the phase space is the product of two cylinders of widths r_1 and r_2) so that generically stability follows from KAM theory.

4 Stability of Hamiltonian Relative Equilibria

In this section we apply the results of Section 3 on the stability of Poisson equilibria to relative equilibria of Hamiltonian systems which are invariant under free group actions. To do this we use the fact that relative equilibria are equilibria of the symmetry reduced dynamics on the Poisson orbit manifold.

Let P be a finite dimensional symplectic manifold with a free symplectic action of a Lie group G . Assume that the action has an Ad_G^* -equivariant momentum map $J : P \rightarrow \mathfrak{g}^*$ with respect to the coadjoint action of G on $\mathfrak{g}^*$. Let H be a G -invariant function on P , X_H the corresponding equivariant Hamiltonian vector field and Φ_t its equivariant flow. The flow preserves the level sets of both H and J . Note that a fibre $J^{-1}(\mu)$ is only G_μ -invariant and the restriction of the flow to the fibre is G_μ -equivariant.

By definition, a point $p_e \in P$ is a relative equilibrium if there exists a *generator* $\xi_e \in \mathfrak{g}$ such that $X_H(p_e) = \xi_e p_e$. This is equivalent to p_e being a critical point of $H_{\xi_e} = H - J_{\xi_e}$ where $J_{\xi_e}(p) = J(p)(\xi_e)$. Note that the trajectory of X_H through p_e is $\exp(t\xi_e)p_e$. Equivariance and conservation of J implies that $\xi_e \in \mathfrak{g}_{\mu_e}$.

Definition 4.1 A relative equilibrium p_e is:

1. *G-stable* if for every G -invariant neighbourhood U of p_e there is a neighbourhood V such that $\Phi_t(p) \in U$ for all $p \in V$ and all t ;
2. *Leafwise stable* if it is G_{μ_e} -stable for the restriction of Φ_t to the momentum level set $J^{-1}(\mu_e)$.

Thus a relative equilibrium is leafwise stable if it is stable (mod G_μ) to momentum preserving perturbations and G -stable if it is stable (mod G) to all perturbations.

For simplicity we restrict the discussion in this paper to free, proper group actions. In [37] we will take advantage of the generality of the topological stability lemma to extend the stability theory to actions with nontrivial isotropy subgroups. For a free, proper action, the orbit space P/G is a smooth manifold which inherits a Poisson structure from the symplectic structure on P . The Hamiltonian H descends to a function h on P/G for which the corresponding Poisson flow is the flow ϕ_t on P/G induced by Φ_t . The orbit $x_e = Gp_e \in P/G$ is an equilibrium point of ϕ_t . Moreover the relative equilibrium p_e is G -stable if and only if x_e is Lyapunov stable in the usual sense, and is leafwise stable in the sense of Definition 4.1 if and only if x_e is leafwise stable in the sense of §3. We can therefore expect to lift the theory developed in §3 to obtain both topological and derivative tests for the G -stability of relative equilibria.

The topological and derivative tests for the G -stability of Hamiltonian relative equilibria are contained in Sections 4.1 and 4.3, respectively. Section 4.2 introduces *transverse Poisson structures* as tools for describing local leafspace topology and uses this to discuss some special cases. At the end of Section 4.3 we discuss applications to relative equilibria of systems that are invariant under actions of the Euclidean groups $SE(2)$ and $SE(3)$ and an example of Libermann and Marle [22].

4.1 G -Stability by Topological Methods

In this section we obtain topological stability criteria for relative equilibria of G -invariant Hamiltonians on P by applying the results of Section 3.1 for Poisson equilibria to the symmetry reduced flow on the Poisson orbit manifold P/G .

The main result of the section is the following theorem. It is a generalisation to non-compact free group actions of Montaldi's stability theorem [31] for compact groups. An extension to general proper actions is given in [37].

Theorem 4.2 *Let H be a G -invariant Hamiltonian on P with a relative equilibrium at p_e . Let S be a slice transverse to the orbit Gp_e at p_e . Then p_e is G -stable if there exists an open neighbourhood U_S of p_e in S and an open neighbourhood U_{μ_e} of μ_e in $\mathfrak{g}^*$ such that*

$$H^{-1}(H(p_e)) \cap J^{-1}(T_2^{U_{\mu_e}}(\mu_e)) \cap U_S = \{p_e\}. \quad (4.4)$$

This remains true if H is replaced by any conserved quantity with values in any Hausdorff space.

The proof of Theorem 4.2 uses the slice S as a local model for the orbit space P/G . The projection from S to P/G induces a Poisson structure on S that is isomorphic to that on the corresponding open neighbourhood of Gp_e in P/G . The following result shows that the T_2 -set of the Poisson structure on S at p_e is just the pullback of the T_2 -set of the Poisson structure on $\mathfrak{g}^*$ at μ_e .

Lemma 4.3 *Let S denote a slice to Gp_e at p_e .*

1. *The symplectic leaves of the induced Poisson structure on S are the connected components of the intersections $J^{-1}(\mathcal{O}) \cap S$, where $\mathcal{O}$ is coadjoint orbit in $\mathfrak{g}^*$.*

2. There exist arbitrarily small neighbourhoods U_S of p_e in S and U_{μ_e} of μ_e in $\mathfrak{g}^*$ such that

$$T_2^{U_S}(p_e) = J^{-1}(T_2^{U_{\mu_e}}(\mu_e)) \cap U_S. \quad (4.5)$$

Proof. The first part follows from the fact that the symplectic leaves of P/G are the connected components of $J^{-1}(\mathcal{O})/G$.

For the second part we note that since J is a submersion there exist arbitrarily small neighbourhoods U_{p_e} of p_e in P and U_{μ_e} of μ_e in $\mathfrak{g}^*$ such that if M is a connected set in U_{μ_e} then $J^{-1}(M) \cap U_{p_e}$ is also connected. If $U_S = U_{p_e} \cap S$ then this also implies that $J^{-1}(M) \cap U_S$ is connected, since this last intersection is homeomorphic to the projection of $J^{-1}(M) \cap U_{p_e}$ to P/G . It follows that the momentum map induces a homeomorphism between the leaf space of U_S and that of U_{μ_e} , which in turn implies (4.5). ■

Proof of Theorem 4.2. We may assume that the neighbourhoods U_S and U_{μ_e} in (4.4) also satisfy the conclusions of Lemma 4.3. The condition (4.4) therefore implies that the restriction $H|_S$ isolates p_e in $T_2^{U_S}(p_e)$ and so Theorem 3.1 implies that p_e is Lyapunov stable for the flow on S generated by $H|_S$ and the induced Poisson structure on S . Identifying S with a neighbourhood of Gp_e in P/G and the flow on S with the quotient of the flow on P generated by H gives the result.

The statement that H can be replaced by any conserved quantity with values in any Hausdorff space follows immediately from the analogous statement in Theorem 3.1. ■

Remark 4.4 It follows from Lemma 4.3 that the leaf space of P/G is Hausdorff at Gp_e if and only if $\mathfrak{g}^*/G$ is Hausdorff at $G\mu_e$. If this holds then the condition (4.4) reduces to

$$H^{-1}(H(p_e)) \cap J^{-1}(\mu_e) \cap U_S = \{p_e\}, \quad (4.6)$$

which means that H isolates $G_{\mu_e}p_e$ in its momentum level set. Sufficient conditions on μ_e for $\mathfrak{g}^*/G$ to be Hausdorff at $G\mu_e$ are given in Proposition 4.10. In general condition (4.6) always implies that p_e is leafwise stable.

4.2 Transverse Poisson Structures and Tame Generators

Weinstein's local splitting theorem for a Poisson manifold X [47, Theorem 2.1] states that any point $x \in X$ has a neighbourhood U that is isomorphic as a Poisson manifold to the product of a neighbourhood of x in the leaf $L(x)$ and a transverse Poisson space Σ . The isomorphism class of this transverse Poisson structure does not depend on either the point x in $L(x)$ or the choice of transverse section Σ . In Section 4.2.1 we give a detailed description of the transverse Poisson space at a point $\mu \in \mathfrak{g}^*$ and use this to give sufficient conditions for $\mathfrak{g}^*/G$ to be locally Hausdorff at $G\mu$ and for a generator $\xi \in \mathfrak{g}_\mu$ to be tame. In Section 4.2.2 we show that the transverse Poisson structure at $Gp \in P/G$ is isomorphic to that at $\mu = J(p)$ and use this to describe the tame generators on P/G at Gp .

4.2.1 Coadjoint Actions

In this section we will define an explicit choice of a transverse Poisson space to the coadjoint orbit $G\mu$ at μ in $\mathfrak{g}^*$ and describe some of its properties.

Let G_μ denote the isotropy subgroup of $\mu \in \mathfrak{g}^*$ and $\mathfrak{g}_\mu$ the isotropy subalgebra. Let $\mathfrak{n}_\mu$ denote a complement to $\mathfrak{g}_\mu$ in $\mathfrak{g}$, so that $\mathfrak{g} = \mathfrak{g}_\mu \oplus \mathfrak{n}_\mu$ and $\mathfrak{g}^* = \mathfrak{g}_\mu^\circ \oplus \mathfrak{n}_\mu^\circ$, where the superscript $^\circ$ denotes an annihilator. The decomposition of $\mathfrak{g}$ induces an isomorphism between $\mathfrak{n}_\mu^\circ$ and $\mathfrak{g}_\mu^*$. An easy calculation shows that $\mathfrak{g}\mu = \mathfrak{g}_\mu^\circ$, so the affine subspace $\mu + \mathfrak{n}_\mu^\circ$ is transverse to the coadjoint orbit $G\mu$ at μ . It then follows from [47, Proposition 4.1] that there is an induced Poisson structure on a neighbourhood Σ of μ in $\mu + \mathfrak{n}_\mu^\circ$. This transverse Poisson structure can be described explicitly as follows [10, 41]. Let $\pi_{\mathfrak{g}_\mu^\circ}$ denote the projection from $\mathfrak{g}^*$ to $\mathfrak{g}_\mu^\circ$ with kernel $\mathfrak{n}_\mu^\circ$. Since Σ is affine, its tangent spaces are all equal to $\mathfrak{n}_\mu^\circ$, which is identified with $\mathfrak{g}_\mu^*$. Thus the derivatives of functions on Σ can be regarded as taking values in $\mathfrak{g}_\mu$.

Proposition 4.5 [10, 41] *There exists a neighbourhood $V \subseteq \mathfrak{n}_\mu^\circ$ of 0 such that:*

1. *For every $\nu \in V$ and every $\xi \in \mathfrak{g}_\mu$ the equation*

$$\pi_{\mathfrak{g}_\mu^\circ}(\text{ad}_{\xi+\eta}^*(\mu + \nu)) = 0 \quad (4.7)$$

has a unique solution $\eta = \eta_{\mu+\nu}(\xi) \in \mathfrak{n}_\mu$ and the map $j_\mu(\nu) : \mathfrak{g}_\mu \rightarrow \mathfrak{g}$ defined by $j_\mu(\nu)\xi = \xi + \eta_{\mu+\nu}(\xi)$ is linear, depends smoothly on ν and satisfies $j_\mu(0)\xi = \xi$.

2. *A Poisson structure is defined on $\Sigma = \mu + V$ by the bracket*

$$\{f, g\}(\mu + \nu) = -\langle \mu + \nu, [j_\mu(\nu)d_\nu f(\mu + \nu), j_\mu(\nu)d_\nu g(\mu + \nu)] \rangle$$

where f, g are smooth functions on Σ and $[\cdot, \cdot]$ is the Lie bracket on $\mathfrak{g}$.

3. *The Poisson structure on a neighbourhood of μ in $\mathfrak{g}^*$ is isomorphic to the product of this Poisson structure on Σ and the Kostant-Kirillov-Souriau symplectic structure on a neighbourhood of μ in $G\mu$.*

The symplectic leaves of the Poisson structure of Σ are the intersections of Σ with the coadjoint orbits on $\mathfrak{g}^*$. Next we characterise these using some notation and results from [49].

Definition 4.6 For each ν in a neighbourhood $V \subseteq \mathfrak{n}_\mu^\circ$ of 0 define $Z_{\mu,\nu}$ to be the connected component containing the identity of

$$\tilde{Z}_{\mu,\nu} \equiv \{g \in G : \text{Ad}_g^*(\mu + \nu) \in \mu + V\},$$

and define $Z_{\mu,V} = \bigcup_{\nu \in V} Z_{\mu,\nu}$. Clearly $Z_{\mu,0} = G_\mu^\circ$ and $Z_{\mu,\nu} \supseteq G_{\mu+\nu}^\circ$.

Proposition 4.7

1. *For each ν sufficiently close to 0 there exists a neighbourhood W of the identity in G such that $Z_{\mu,\nu} \cap W$ is a manifold of dimension $\dim G_\mu$. Its tangent space at the identity is $T_{\text{id}}Z_{\mu,\nu} = j_\mu(\nu)\mathfrak{g}_\mu$.*
2. *For V sufficiently small the symplectic leaf through $\mu + \nu$ of the transverse Poisson structure on $\Sigma = \mu + V$ is equal to $Z_{\mu,\nu}(\mu + \nu) \cap \Sigma$.*

The second statement says essentially that the symplectic leaves of Σ are the ‘orbits’ of $Z_{\mu,V}$. However in general $Z_{\mu,\nu}$ and $Z_{\mu,V}$ are not groups.

Proof. The first statement was proved in [49] and follows from an application of the implicit function theorem. The second statement is an immediate consequence of Definition 4.6 and the fact that the symplectic leaves of Σ are its intersections with the coadjoint orbits of G . ■

The set of generators (see Definition 3.5) at μ for the Poisson structure on $\mathfrak{g}^*$ is $(\mathfrak{g}\mu)^\circ = \mathfrak{g}_\mu$. Since any element of $T_\mu^*\Sigma$ is a generator for the Poisson structure on Σ , and $T_\mu^*\Sigma = \mathfrak{n}_\mu^* = (\mathfrak{g}\mu)^\circ = \mathfrak{g}_\mu$, the generators for the two Poisson structures coincide.

By Weinstein's local splitting theorem, for an appropriate neighbourhood U_μ of $\mu \in \mathfrak{g}^*$, the set $T_2^{U_\mu}(\mu)$ is the product of $T_2^\Sigma(\mu)$ with a neighbourhood of μ in the leaf $G\mu$. Thus, a generator $\xi \in \mathfrak{g}_\mu$ is *tame* as a generator on $\mathfrak{g}^*$ if and only if it is tame as a generator on Σ , ie if and only if the equivalent conditions $T_\mu(T_2^\Sigma(\mu)) \subseteq \text{ann}_{\mathfrak{g}_\mu^*} \xi$ and $T_\mu(T_2^{U_\mu}(\mu)) \subseteq \text{ann}_{\mathfrak{g}^*} \xi$ hold. The annihilator of ξ in $\mathfrak{g}_\mu^*$ is denoted by $\text{ann}_{\mathfrak{g}_\mu^*} \xi$, and other annihilators in a similar way. Note that the property of being tame does not depend on the choice of the transverse section Σ .

In the next section we will need a stronger form of tameness.

Definition 4.8 A generator $\xi \in T_\mu^*\Sigma \cong \mathfrak{g}_\mu$ is *very tame* if $T_2^\Sigma(\mu) \subseteq \mu + \text{ann} \xi$.

Clearly very tame generators are always tame, but in general the converse does not hold, see Remark 4.11.

There are two important special cases of transverse Poisson structures.

Definition 4.9 We say that $\mu \in \mathfrak{g}^*$ is:

1. *Regular* if $\dim \mathfrak{g}_\nu = \dim \mathfrak{g}_\mu$ for every ν in a neighbourhood of μ .
2. *Split* if there exists a G_μ° -invariant complement $\mathfrak{n}_\mu$ to $\mathfrak{g}_\mu$ in $\mathfrak{g}$.

Here G° denotes the identity component of G . Note that μ is regular if and only if it is a regular point for the Lie-Poisson structure on $\mathfrak{g}^*$ in the sense of Section 3.2, since the symplectic leaves are the coadjoint orbits and the definition above implies that these have constant dimension near regular points.

It is clear that if there exists a G -invariant inner product on $\mathfrak{g}$, as in the case of compact or Abelian groups, then every μ is split, the T_2 -sets are trivial and every generator is very tame. Some other results are given in the following proposition.

Proposition 4.10

1. If μ is regular then $\mathfrak{g}_\mu$ is Abelian, $Z_{\mu,\nu} = G_{\mu+\nu}^\circ$ for ν small, the transverse Poisson structure on Σ is trivial, $\mathfrak{g}^*/G$ is Hausdorff at $G\mu$ and every generator is very tame. The set of regular points in $\mathfrak{g}^*$ is open and dense.
2. Let μ be split. Then $Z_{\mu,\nu} = G_\mu^\circ$ for every ν , the transverse Poisson structure on Σ is isomorphic to the standard Lie-Poisson structure on $\mathfrak{g}_\mu^*$ and its symplectic leaves are the coadjoint orbits of G_μ° on $\mathfrak{g}_\mu^*$. Moreover a generator $\xi \in \mathfrak{g}_\mu$ is tame if and only if it is very tame, and a sufficient condition for $\xi \in \mathfrak{g}_\mu$ to be tame is that its adjoint orbit $G_\mu^\circ \xi$ is bounded. Finally $\mathfrak{g}^*/G$ is Hausdorff at $G\mu$ if there is a G_μ° -invariant inner product on $\mathfrak{g}_\mu$ and in particular if G_μ is compact.

Proof. The results for regular μ are due essentially to Lie [23] and Duflo and Vergne [11]. For split μ the results in the first sentence of Part 2 follow from Proposition 4.5. See also [47, 10] and references therein. For the remaining statements of Part 2 note that since μ is split we can assume that $\mu = 0$, $G_\mu = G$ and Σ is an open neighbourhood of 0 in $\mathfrak{g}^*$. The linearity of the action of G on $\mathfrak{g}^*$ implies that if $\nu \in T_2^\Sigma(0)$ then $\lambda\nu \in T_2^\Sigma(0)$ for all $\lambda \in \mathbb{R}$. It follows that $\nu \in T_0(T_2^\Sigma(0))$ and so $T_2^\Sigma(0) \subseteq T_0(T_2^\Sigma(0))$. Hence, if ξ annihilates $T_0(T_2^\Sigma(0))$ then it annihilates $T_2^\Sigma(0)$, and so the sets of tame and very tame generators coincide.

If $\nu \in T_2^\Sigma(0)$ then there are sequences $\nu_n \in \Sigma$ and $g_n \in G_\mu^\circ$, such that $\nu_n \rightarrow \nu$ and $g_n\nu_n \rightarrow 0$ as $n \rightarrow \infty$. We need to show that $\langle \nu, \xi \rangle = 0$ if $G^\circ\xi$ is bounded. But $\langle \nu, \xi \rangle = \lim_{n \rightarrow \infty} \langle \nu_n, \xi \rangle = \lim_{n \rightarrow \infty} \langle g_n\nu_n, g_n\xi \rangle = 0$ since $g_n\nu_n \rightarrow 0$, $g_n\xi \in G^\circ\xi$ and $G^\circ\xi$ is bounded. ■

Remark 4.11 In general regular momenta need not be split and their $Z_{\mu,\nu}$ need not be equal to G_μ° . Examples include nonzero points on the nilpotent cone of $sl(2, \mathbb{R})^*$ (see Example 3.10 and Example 4.4 of [41]). If μ is non-split and non-regular then there may be tame generators $\xi \in \mathfrak{g}_\mu$ which are not very tame. In Example 3.15, where μ is a subregular nilpotent element of $sl(3, \mathbb{R})^*$, the fact that $T_\mu(T_2^{U\Sigma}(\mu))$ is one dimensional while $T_2^{U\Sigma}(\mu)$ is two dimensional implies that there exist generators ξ which annihilate $T_\mu(T_2^{U\Sigma}(\mu))$ but not $T_2^{U\Sigma}(\mu)$ itself.

4.2.2 The Orbit Space P/G

We now give a description of how the transverse Poisson structure at $\mu \in \mathfrak{g}^*$ is related to local Poisson structure of P/G near Gp .

As in Section 4.1 we identify a neighbourhood of Gp in P/G with a slice S transverse to the orbit Gp in P . By Lemma 4.3 the symplectic leaf $L^S(p)$ of the Poisson structure on S is the connected component of $J^{-1}(G\mu) \cap S$ containing p . Since J is a submersion the image $\Sigma = J(S)$ is a submanifold of $\mathfrak{g}^*$ through μ that is transverse to $G\mu$ and so has an induced transverse Poisson structure. The fact that J is a Poisson map [26] implies that the transverse Poisson space at $p \in S$ is isomorphic to that at $\mu \in \mathfrak{g}^*$. It follows from Weinstein's local splitting theorem that a sufficiently small neighbourhood of p in S is isomorphic as a Poisson manifold to a neighbourhood of (p, μ) in the product $L^S(p) \times \Sigma$.

Recall that p_e is a relative equilibrium with generator ξ_e if and only if it is a critical point of H_{ξ_e} . The derivative tests which we give in Section 4.3 state that if the generator is tame (or very tame) and the Hessian $d^2H_{\xi_e}(p_e)$ is definite when restricted to certain subspaces of the normal space N to Gp_e at p_e , then p_e is G -stable. The *Witt decomposition* [6] of the normal space to any group orbit Gp splits it into two components $N = N_0 \oplus N_1$, where N_1 is the *symplectic normal space*, ie a maximal subspace of N for which the restriction of the symplectic form on T_pP is nondegenerate. More explicitly N_1 is a complement to $T_p(G_\mu p)$ in $\ker dJ(p)$. For free group actions it is isomorphic to the tangent space at Gp to the leaf $L(Gp)$ through Gp in P/G , ie the Marsden-Weinstein reduced phase space [28].

The derivative $dJ(p_e)$ maps a complement N_0 to N_1 in N isomorphically to $T_{\mu_e}\Sigma$, the normal space to $G\mu$ at μ and hence identifies it with $\mathfrak{n}_\mu^\circ \cong \mathfrak{g}_\mu^*$. Using this identification it can be endowed with the transverse Poisson structure at μ described in Section 4.2.1. The normal space N can then be given the Poisson structure obtained by taking the product of this structure on N_0 with the symplectic structure on N_1 .

The following result is a version of Weinstein's local splitting of P/G , or equivalently a slice S , due to Guillemin, Sternberg and Marle.

Theorem 4.12 [12, 25] *Let S be a slice in P transverse to Gp at p such that $J(S) \subset \Sigma = \mu + \mathfrak{n}_\mu^\circ$. Then there exists a Poisson isomorphism from S to an open neighbourhood of the origin in $N = N_0 \oplus N_1$ with its product Poisson structure such that the restriction of the momentum map J to S is given by $J_S(\nu, w) = \mu + \nu$ where $\nu \in N_0 \cong \mathfrak{n}_\mu^\circ$ and $w \in N_1$.*

It follows from the local decomposition of P/G given in Theorem 4.12 that the set of generators at Gp_e can also be identified with $T_\mu^*\Sigma$, and hence with $\mathfrak{g}_\mu$, and that a generator is (very) tame for the Poisson structure on P/G if and only if under this identification it is (very) tame for the transverse Poisson structure on $\Sigma \cong \mathfrak{g}_\mu^*$.

4.3 T_2 -Energy-Momentum-Casimir Method

We now turn to the task of lifting the derivative tests of Section 3.2 from P/G to the symplectic phase space P . Our main results are Theorem 4.15 and Corollary 4.17 which, when combined with the results of Section 5, generalise all previous published results for free group actions. Their extensions to general proper actions will be given in [37].

Before giving our main result we present a simple corollary of Theorem 4.2. This will be generalised by Corollary 4.17.

Corollary 4.13 *Let p_e be a relative equilibrium of H with generator ξ_e . Suppose that $\mathfrak{g}^*/G$ is Hausdorff at $\mu_e = J(p_e)$ and that the Hessian $d^2H_{\xi_e}(p_e)$ is (positive or negative) definite when restricted to any symplectic normal space at p_e . Then p_e is G -stable.*

If $\mathfrak{g}^*/G$ is not Hausdorff at μ_e then the definiteness of $d^2H_{\xi_e}(p_e)$ on a symplectic normal space only implies that p_e is leafwise stable.

By Proposition 4.10 the space $\mathfrak{g}^*/G$ is Hausdorff at μ_e if either μ_e is regular or μ_e is split and there exists a G_{μ_e} -invariant inner product on $\mathfrak{g}_{\mu_e}^*$. The G -stability result for μ_e regular is Theorem 8.17 of Chapter IV of [22]. The condition that μ_e is split and there exists a G_{μ_e} -invariant inner product on $\mathfrak{g}_{\mu_e}^*$ is implied by the existence of a G_{μ_e} -invariant inner product on the whole of $\mathfrak{g}^*$, which is always true for compact groups. Under this stronger hypothesis [19, 33] show that definiteness of $d^2H_{\xi_e}(p_e)$ on a symplectic normal space implies that p_e is actually G_{μ_e} -stable, generalising a result of [35] for compact groups G . We recover this result by combining Corollary 4.13 with Corollary 5.4 in Section 5.

Proof. If S is a slice to Gp_e at p_e the symplectic normal space can be identified with $T_{p_e}(J^{-1}(\mu_e) \cap S)$ and the definite Hessian, together with the fact that p_e is a critical point of H_{ξ_e} , implies that p_e is isolated in $H_{\xi_e}^{-1}(H_{\xi_e}(p_e)) \cap J^{-1}(\mu_e) \cap S$. Since J_{ξ_e} is constant on $J^{-1}(\mu_e)$ this implies that p_e is also isolated in $H^{-1}(H(p_e)) \cap J^{-1}(\mu_e) \cap S$. If $\mathfrak{g}^*/G$ is Hausdorff at μ_e then this is equivalent to condition (4.4) of Theorem 4.2 and so p_e is also G -stable. ■

In Theorem 4.15 below we generalise the ‘energy-momentum method’ of Corollary 4.13 to obtain criteria for the G -stability of Hamiltonian relative equilibria for general non-compact symmetry groups and non-regular momentum values. Before formulating the theorem we make some preliminary remarks on smoothings and Casimirs. A *local Casimir* is a continuous function $C : U_{\mu_e} \rightarrow \mathbb{R}$ on a neighbourhood U_{μ_e} of μ_e which is constant on the connected components of the coadjoint orbits intersected with U_{μ_e} .

Remark 4.14

1. It follows from Lemma 4.3 that for a sufficiently small slice S to Gp_e at p_e we have $T_2^S(p_e) = J^{-1}(T_2^\Sigma(\mu_e)) \cap S$ where $\Sigma = J(S)$. Since J is a submersion at p_e this means that $\{B_i\}_{i=1}^n$ is a smoothing of $T_2^S(p_e)$ if and only if $\{J(B_i)\}_{i=1}^n$ is a smoothing of $T_2^\Sigma(\mu_e)$.
2. Every local Casimir $C : U_{\mu_e} \rightarrow \mathbb{R}$ with $\Sigma \subset U_{\mu_e}$ is constant on $T_2^\Sigma(\mu_e)$. By Lemma 4.3 the function $C \circ J$ is a Casimir on every slice S satisfying $J(S) \subseteq U_{\mu_e}$. Consequently, as in Section 3, if $\{B_i\}_{i=1}^n$ is a smoothing of $T_2^S(p_e)$ the function $C \circ J|_{B_i}$ has a critical point at p_e for each i .
3. If $\{B_i\}_{i=1}^n$ is a smoothing of $T_2^S(p_e)$ a relative equilibrium p_e with generator $\xi_e \in \mathfrak{g}_{\mu_e}$ is a critical point of $H|_{B_i}$ for each i if and only if ξ_e is tame. To see this, note that $dH(p_e) = \langle dJ(p_e), \xi_e \rangle$ annihilates $T_{p_e} T_2^S(p_e) = dJ(p_e)^{-1}(T_{\mu_e} T_2^\Sigma(\mu_e))$ if and only if ξ_e annihilates $T_{\mu_e}(T_2^\Sigma(\mu_e))$.

The next theorem provides two Hessian tests for the G -stability of relative equilibria with tame generators. The first uses the second derivatives of the restrictions of H to the manifolds B_i in a smoothing of $T_2^S(p_e) = J^{-1}(T_2^\Sigma(\mu_e)) \cap S$. In general it is not possible to calculate the second derivative of $H|_S$ and then restrict it to the tangent spaces $T_{p_e} B_i$ because H need not have a critical point at p_e . If $J^{-1}(T_2^\Sigma(\mu_e)) = J^{-1}(\mu_e)$ then this can be circumvented by replacing the second derivative of H by that of $H_{\xi_e} = H - J_{\xi_e}$, which does have a critical point at p_e . This is valid because J_{ξ_e} is constant on this level set of J . We used this idea in the proof of Corollary 4.13.

If the set $J^{-1}(T_2^\Sigma(\mu_e))$ is larger than the level set of J then it may not be possible to replace H by H_{ξ_e} . However if ξ_e is *very tame*, as defined in Definition 4.8, then J_{ξ_e} is constant on the whole of $J^{-1}(T_2^\Sigma(\mu_e))$ and it is possible to proceed as in the Hausdorff case. This gives the second test. Both tests can be strengthened by the addition of Casimirs, as stated in the theorem.

Theorem 4.15 (*T₂-Energy-Momentum-Casimir Method*) *Let p_e be a relative equilibrium of H with generator ξ_e . Let S be a slice to Gp_e at p_e and $\{B_i\}_{i=1}^n$ a smoothing of $T_2^S(p_e)$.*

1. *If ξ_e is a tame generator then p_e is G -stable if for each $1 \leq i \leq n$ there is a smooth local Casimir C_i such that the Hessian $d^2((H + C_i \circ J)|_{B_i})(p_e)$ is definite on $T_{p_e} B_i$.*
2. *If ξ_e is a very tame generator for the transverse Poisson structure on $\Sigma = J(S)$ then p_e is G -stable if for each $1 \leq i \leq n$ there is a smooth local Casimir C_i with a critical point at μ_e and a Hessian $d^2(H_{\xi_e} + C_i \circ J)(p_e)$ that is definite when restricted to $T_{p_e} B_i$.*

Proof.

1. By Remark 4.14.2 the functions $C_i \circ J$ are Casimirs on S and so 1. follows from Theorem 3.6.

2. If ξ_e is very tame then for all $p \in T_2^S(p_e) = J^{-1}(T_2^\Sigma(\mu_e))$ we have:

$$J_{\xi_e}(p) = \langle J(p), \xi_e \rangle = \langle J(p) - \mu_e, \xi_e \rangle + \langle \mu_e, \xi_e \rangle$$

since $J(p) \in T_2^\Sigma(\mu_e) \subseteq \mu_e + \text{ann } \xi_e$. Let $f_i = H|_S + C_i \circ J|_S$, $\tilde{f}_i = H_{\xi_e}|_S + C_i \circ J|_S$, $f = (f_1, \dots, f_n)$ and $\tilde{f} = (\tilde{f}_1, \dots, \tilde{f}_n)$. Then on $J^{-1}(T_2^\Sigma(\mu_e))$ the restrictions of f and $\tilde{f}$ differ by a constant vector.

If C_i has a critical point at μ_e then $\tilde{f}_i$ has a critical point at p_e and

$$d^2 \tilde{f}_i(p_e)|_{T_{p_e} B_i} = d^2(\tilde{f}_i|_{B_i})(p_e)$$

It follows that if the restriction of the Hessian of $\tilde{f}_i$ to $T_{p_e} B_i$ is definite then p_e is an isolated point in $(\tilde{f}_i|_{B_i})^{-1}(\tilde{f}_i(p_e))$ for each i and so an isolated point in $\tilde{f}^{-1}(\tilde{f}(p_e)) \cap J^{-1}(T_2^\Sigma(\mu_e))$. Since on $J^{-1}(T_2^\Sigma(\mu_e))$ the functions f and $\tilde{f}$ differ only by a constant this implies that p_e is an isolated point of $f^{-1}(f(p_e)) \cap J^{-1}(T_2^\Sigma(\mu_e))$. The proof now follows from Theorem 4.2. ■

We will extend the T_2 -Energy-Momentum-Casimir Method to general proper actions in [37].

Remark 4.16

1. By Remark 4.14.1 inclusion of Casimirs in the theorem is unnecessary if the smoothing $\{J(B_i)\}_{i=1}^n$ of $T_2^\Sigma(\mu_e)$ satisfies conditions similar to those stated at the end of §3.2. In particular Casimirs are unnecessary if $T_2^\Sigma(\mu_e)$ is a manifold, and the inclusion of C_i is unnecessary for any i such that (3.1) holds for the Poisson space Σ .
2. If $T_2^S(p_e)$ is a one dimensional submanifold of S then it is not difficult to formulate a derivative test for the G -stability of p_e analogous to that of Corollary 3.2 for Poisson equilibria.
3. Theorem 4.15 can be extended to weak smoothings, as in Remark 3.7.

We announced the following corollary of Theorem 4.15 in [50, Theorem 2]. Although the result is not always optimal, it gives a generalisation of the energy-momentum method of [19, 35, 33] to non-compact symmetry groups that, as will be seen in Corollaries 4.18 and 4.19, is optimal for the Euclidean symmetry groups that are most likely to arise in applications. Recall from Section 4.2.2 that the normal space N to Gp_e at p_e decomposes as $N_0 \oplus N_1$ where N_1 is a symplectic normal space and $N_0 \cong \mathfrak{g}_\mu^*$.

Corollary 4.17 (Simple Energy-Momentum Method) *Let $\mathfrak{t}_{\mu_e} \subseteq \mathfrak{g}_{\mu_e}$ be the space of very tame generators at μ_e and $\mathfrak{w}_{\mu_e}^* = \text{ann}_{\mathfrak{g}_{\mu_e}^*} \mathfrak{t}_{\mu_e}$. Then the relative equilibrium p_e with momentum μ_e is G -stable if its generator ξ_e is very tame and $d^2 H_{\xi_e}(p_e)|_{\mathfrak{w}_{\mu_e}^* \oplus N_1}$ is definite.*

We call the subspace $\mathfrak{w}_{\mu_e}^* = \text{ann}_{\mathfrak{g}_{\mu_e}^*} \mathfrak{t}_{\mu_e}$ of $\mathfrak{g}_{\mu_e}^*$ the space of *wild momenta*.

Proof. Choose the slice S at p_e as in Theorem 4.12 and identify it with the normal space $N = N_0 \oplus N_1$. By definition very tame generators annihilate the set $T_2^\Sigma(\mu_e)$ and so the space $B = \mathfrak{w}_{\mu_e}^* \oplus N_1 \subseteq N$ is a weak smoothing of $T_2^S(p_e)$. The corollary therefore follows from Remark 4.16.3. ■

For the case of semidirect products of compact groups and vector spaces Leonard and Marsden [18] also identified an intermediate set between N_1 and $T_p P$ on which definiteness of $d^2 H_{\xi_e}(p_e)$ implies stability. Corollary 4.17 generalises their results.

In [50] we restricted to the case where μ_e is split and *defined* generators $\xi \in \mathfrak{g}_{\mu_e}$ to be tame if $G_{\mu_e}^o \xi$ is bounded. By Proposition 4.10.2 this definition of tame generators is stronger than the notion of very tame generators used in this paper.

We now show how the energy-momentum method of this section applies to relative equilibria of Euclidean invariant Hamiltonian systems. We treat the cases where the symmetry

group is the special Euclidean group $G = SE(2) = SO(2) \ltimes \mathbb{R}^2$ of rotations and translations of the plane and the special Euclidean group $G = SE(3) = SO(3) \ltimes \mathbb{R}^3$ of three-space. Relative equilibria of Euclidean invariant Hamiltonian systems have been studied for several systems, including the dynamics of underwater vehicles [17, 18] and systems of point vortices [32], and the statics of elastic rods [30].

In the case of $SE(2)$ the coadjoint action is

$$(R, a) \cdot (\mu^r, \mu^a) = (\mu^r + R\mathbb{J}\mu^a \cdot a, R\mu^a)$$

where we have represented elements of $SO(2)$ as 2×2 matrices and $\mathbb{J} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. From this one sees that $(\mu^r, \mu^a) \mapsto |\mu^a|$ is a Casimir, and that the coadjoint orbits are the cylinders about the μ^r axis together with the points on the μ^r axis. Consequently $\mu \in se(2)^*$ is regular if $\mu^a \neq 0$ in which case $G_\mu = \{(1, t\mu_e^a) : t \in \mathbb{R}\}$, and $\mu \in se(2)^*$ is non-regular if $\mu^a = 0$ in which case $G_\mu = SE(2)$. Any regular momentum (μ^r, μ^a) is split since the transverse section $\{(\mu^r, t\mu^a) : t \in \mathbb{R}\}$ is G_μ invariant, while the non-regular momenta are trivially split since they are at zero dimensional coadjoint orbits. Any two neighbourhoods of two non-regular momenta both meet sufficiently narrow cylinders about the μ^r -axis, so if μ is non-regular then $T_2(\mu)$ is the μ^r -axis. Given a non-regular momentum (μ^r, μ^a) , a generator (ξ^r, ξ^a) is tame if it annihilates the tangent space to this T_2 -set or equivalently if and only if $\xi^r = 0$, and every tame generator is very tame. Casimirs are not required for an application of Theorem 4.15 in this case since the T_2 -sets are manifolds. The T_2 -set at the nonregular momenta is equal to the annihilator of the (very) tame generators and so Theorem 4.15 is equivalent to Corollary 4.17 here. Summarising all this gives the following corollary.

Corollary 4.18 *Assume $G = SE(2)$. Let p_e be a relative equilibrium of H with generator $\xi_e = (\xi_e^r, \xi_e^a)$ and momentum $\mu_e = (\mu_e^r, \mu_e^a)$.*

1. *If $\mu_e^a \neq 0$ then μ_e is regular, $G_{\mu_e} = \{(1, t\mu_e^a) : t \in \mathbb{R}\}$, and p_e is G -stable if $d^2H_{\xi_e}(p_e)$ is definite on any complement to $\mathfrak{g}_{\mu_e p_e}$ in $\ker dJ(p_e)$ (ie definite on any symplectic normal space).*
2. *If $\mu_e^a = 0$ then μ_e is non-regular, $G_{\mu_e} = SE(2)$, ξ_e is tame if and only if $\xi_e^r = 0$, and p_e is G -stable if ξ_e is tame and $d^2H_{\xi_e}(p_e)$ is definite on any complement to $\mathfrak{g}p_e$ in $dJ^{-1}(p_e)(so(2)^*)$.*

The coadjoint action of $SE(3)$ is explicitly computed in [26] and the analysis proceeds in a similar way to the $SE(2)$ case. Again the T_2 -sets are subspaces of $se(3)^*$, so that every tame generator is also very tame, even though the momenta which are nonzero and non-regular are not split. The result is the following corollary.

Corollary 4.19 *Assume $G = SE(3)$. Let p_e be a relative equilibrium of H with generator $\xi_e = (\xi_e^r, \xi_e^a)$ and momentum $\mu_e = (\mu_e^r, \mu_e^a)$.*

1. *If $\mu_e^a \neq 0$ then μ_e is regular, $G_{\mu_e} = \{(R, t\mu_e^a) : R\mu_e^a = \mu_e^a, t \in \mathbb{R}\} \cong SO(2) \times \mathbb{R}$, and p_e is G -stable if $d^2H_{\xi_e}(p_e)$ is definite on any complement to $\mathfrak{g}_{\mu_e p_e}$ in $\ker dJ(p_e)$ (ie definite on any symplectic normal space).*

2. If $\mu_e^a = 0$ then μ_e is non-regular,

$$G_{\mu_e} = \begin{cases} \{ (R, a) : R\mu_e^r = \mu_e^r, a \in \mathbb{R}^3 \} \cong SO(2) \ltimes \mathbb{R}^3 & \mu_e^r \neq 0, \\ SE(3) & \mu_e^r = 0, \end{cases}$$

ξ_e is tame if and only if $\xi_e^r = 0$, and p_e is G -stable if ξ_e is tame and $d^2H_{\xi_e}(p_e)$ is definite on any complement to $\mathfrak{g}_{p_e}$ in $dJ^{-1}(p_e)(so(3)^*)$.

For both the $SE(2)$ and $SE(3)$ symmetry groups our theory indicates that, in the case of zero translational momentum, only for purely translating relative equilibria of $SE(2)$ invariant systems is G -stability accessible by energy-momentum methods, since only then is the generator tame. For spinning relative equilibria which have zero translational momentum G -stability may be accessible in low dimensional phase space by a blow up argument coupled with KAM theory, as we will show in [38].

Example 4.20 This example, which is due to Libermann, Marle, and Krishnaprasad, is Exercise (15.10) on page 274 of [22] and is worked out in detail in the appendix of [15]. It was presented as an example of a relative equilibrium which is leafwise stable, but not G -stable. Here we show how it relates to the theory of this paper.

The group G is a semidirect product $G = \mathbb{R} \ltimes \mathbb{R}$ with multiplication

$$(a, b)(a', b') = (a + a', b + e^a b')$$

The phase space P is $T^*G = \mathbb{R}^4 = \{(q^1, q^2, p_1, p_2)\}$ with its standard symplectic form $dq^1 \wedge dp_1 + dq^2 \wedge dp_2$. The action of G on P is the cotangent lift of the left translation action of G on itself:

$$(a, b)(q^1, q^2, p_1, p_2) = (a + q^1, b + e^a q^2, p_1, e^{-a} p_2).$$

The Lie algebra dual is $\mathfrak{g}^* \cong \mathbb{R}^2 = \{(\nu_1, \nu_2)\}$, the coadjoint action is

$$\text{Ad}_{(a,b)^{-1}}^* \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} = \begin{pmatrix} 1 & e^{-a}b \\ 0 & e^{-a} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix},$$

and the Ad^* -equivariant momentum mapping is

$$J(q^1, q^2, p_1, p_2) = \begin{pmatrix} p_1 + p_2 q^2 \\ p_2 \end{pmatrix}.$$

As is evident from the formula for the coadjoint action, there are zero dimensional coadjoint orbits along the ν_1 axis, and the half planes $\nu_2 > 0$ and $\nu_2 < 0$ are two dimensional coadjoint orbits of regular momenta. The T_2 -sets for the non-regular momenta on the ν_2 axis are $\mathfrak{g}^*$, and consequently a relative equilibrium at such a momentum is tame if and only if its generator is zero (ie if and only if it is an equilibrium).

The Hamiltonian considered in [22, 15] is $H = p_2 e^{q^1}$, for which the flow on P is

$$q^1(t) = q^1(0), \quad q^2(t) = q^2(0) + t e^{q^1(0)}, \quad p_1(t) = p_1(0) - t p_2(0) e^{q^1(0)}, \quad p_2(t) = p_2(0).$$

The relative equilibria are given by solving $dH = dJ_{(\dot{a}, \dot{b})}$, which directly leads to the equations

$$p_2 e^{q^1} = 0, \quad 0 = \dot{a} p_2, \quad 0 = \dot{a}, \quad e^{q^1} = \dot{a} q^2 + \dot{b},$$

with solution

$$p_2 = 0, \quad \dot{a} = 0, \quad \dot{b} = e^{q^1}.$$

The momenta of these relative equilibria all have $\nu_2 = 0$ and hence are all non-regular, and the generators all have $\dot{b} \neq 0$, and hence are all wild, so we do not expect that any of the relative equilibria are G -stable. On the other hand the momentum level sets of the non-regular momenta are all group orbits so the Marsden-Weinstein reduced spaces corresponding to those momenta are all points, and consequently all the relative equilibria are leafwise stable.

To explicitly see that the relative equilibria are not G -stable, it is best to examine the flow on the Poisson space P/G . The standard Lie-Poisson reduction to $\mathfrak{g}^*$ is realized by the quotient map given by the left trivialisation $\nu = (\nu_1, \nu_2) = g^{-1}(p_1, p_2) = (p_1, p_2 e^{q^1})$. The flow descends to

$$\nu_1(t) = \nu_1(0) - t\nu_2(0), \quad \nu_2(t) = \nu_2(0)$$

and the relative equilibria descend to the equilibria $\nu_2 = 0$. These equilibria are not stable by direct inspection of the reduced flow: perturbations into $\nu_2 \neq 0$ of such equilibria are carried far from their origin by translation parallel to the ν_1 axis.

5 Beyond G-Stability

In this section we show that G -stable relative equilibria typically satisfy a stronger stability property. To motivate this, suppose $p(t)$ is an integral curve of X_H starting at $p(0) \approx p_e$. If p_e is G -stable then there is a curve $g(t)$ in G such that $g(t)^{-1}p(t) \approx p_e$. By continuity and by equivariance and conservation of the momentum J we have:

$$\mu_e = J(p_e) \approx J(g(t)^{-1}p(t)) = \text{Ad}_{g(t)}^* J(p(t)) = \text{Ad}_{g(t)}^* J(p(0)) \approx \text{Ad}_{g(t)}^* \mu_e.$$

In other words *conservation of momentum ought to imply that trajectories which start close to a G -stable relative equilibrium p_e should remain close to the G_{μ_e} orbit through p_e , and not just the G orbit*. Results of this type on G_{μ_e} -stability, all of which assume Hessian conditions and compactness-related invariant inner products or norms on $\mathfrak{g}$, first appeared in [35], with extensions to non-free actions in [19, 29, 33]. That G -stable relative equilibria are not generally G_{μ_e} -stable in the non-compact case was noticed in [35] and then in [18]. In this paper, assuming only the G -stability of a relative equilibrium, we obtain results on stability which lie between G_{μ_e} -stability and G -stability. The results are proved for free actions, but extend to general proper actions [37].

In Section 5.1 we give the main general theorem. This is specialised to split momenta in Section 5.2, and then applied to Euclidean invariant Hamiltonian systems in Section 5.3.

5.1 A-Stability

We first define a very general stability property for equivariant flows.

Definition 5.1 Let P be a topological space with an action of a group G and a G -equivariant flow $\Phi_t: P \rightarrow P$. For any subset $A \subseteq G$, a point $p_e \in P$ is said to be *A-stable* with respect to the flow Φ_t if for any open neighbourhood $U \subseteq P$ of p_e there exists an open neighbourhood $U' \subseteq P$ of p_e such that if $p \in U'$ then $\phi_t(p) \in AU$ for all $t \in \mathbb{R}$.

If $A \subseteq B \subseteq G$ then if p_e is *A-stable* it is also *B-stable*. It is easy to show that a point p_e is *A-stable* if it is *AW-stable* for any neighbourhood W of $1 \in G$.

Let $\Sigma \subseteq \mu_e + \mathfrak{n}_{\mu_e}^o$ be a transverse section through $G\mu_e$ at $\mu_e = J(p_e)$. For any small neighbourhood V of 0 in $\mathfrak{n}_{\mu_e}^o$ and any point $\nu \in V$ let $Z_{\mu_e, \nu}$ and $Z_{\mu_e, V}$ be the subsets of G as in Definition 4.6. For any neighbourhood W of 1 in G define

$$A_{V,W}(\mu_e) = \bigcup_{g \in W} gZ_{\mu_e, V}g^{-1}.$$

Since $Z_{\mu_e, 0} = G_{\mu_e}^o$ we have $G_{\mu_e}^o \subseteq A_{V,W}(\mu_e) \subseteq G$ for any pair of neighbourhoods V and W .

Theorem 5.2 *If p_e is a G -stable relative equilibrium then p_e is an $A_{V,W}(\mu_e)$ -stable equilibrium for every pair of neighbourhoods V of 0 in $\mathfrak{n}_{\mu_e}^o$ and W of 1 in G .*

Proof. Let S be a slice at p_e which is mapped by J to the transverse section $\Sigma \subseteq \mu_e + \mathfrak{n}_{\mu_e}^o$ at μ_e . By Theorem 4.12 the slice S can be identified with the product of open neighbourhoods of $\{0\}$ in N_0 and N_1 , and in these coordinates the restriction of J to S is given by $J_S(\nu, w) = \mu_e + \nu$ where $\nu \in N_0 = \mathfrak{n}_{\mu_e}^o$. By equivariance a G -invariant neighbourhood of Gp_e can be parametrised by $G \times S$ and in these coordinates the momentum map is given by $J(g, \nu, w) = \text{Ad}_{g^{-1}}^*(\mu_e + \nu)$.

Let Φ_t denote the G -equivariant flow on P and ϕ_t the induced flow on P/G . As in Section 4.1, we identify S with an open neighbourhood of Gp_e in P/G and denote the flow on S by ϕ_t . Let U be a sufficiently small neighbourhood of p_e . We then have $W \times (\mu_e + U_0) \times U_1 \subseteq U$ where W is a neighbourhood of $1 \in G$, and U_0 and U_1 are neighbourhoods of $\{0\}$ in N_0 and N_1 , respectively.

Since Gp_e is Lyapunov stable for the flow ϕ_t on P/G there exists a neighbourhood U'_S of p_e in $U_S = (\mu_e + U_0) \times U_1$ such that $\phi_t(U'_S) \subseteq U_S$ for all t . We may choose U'_S to be a product $(\mu_e + U'_0) \times U'_1$ where U'_0 and U'_1 are again neighbourhoods of $\{0\}$ in N_0 and N_1 , respectively. Since S is a slice, for $g_0 \in G$, $\nu_0 \in U'_0$ and $w_0 \in U'_1$ we have

$$\Phi_t(g_0, \nu_0, w_0) \equiv (g(t), \nu(t), w(t)) = g(t)(1, \nu(t), w(t)) \equiv g(t)\phi_t(\nu_0, w_0).$$

The momentum map J is preserved by the flow Φ_t and so $\text{Ad}_{g(t)^{-1}}^*(\mu_e + \nu(t)) = \text{Ad}_{g_0^{-1}}^*(\mu_e + \nu_0)$ for all t , and hence $\text{Ad}_{g_0^{-1}g(t)}^*(\mu_e + \nu_0) = \mu_e + \nu(t) \in \mu_e + U_0$. If the neighbourhood $U_0 \subseteq V$ is chosen small enough this implies that $g(t) \in g_0Z_{\mu_e, \nu_0}$ for all t (see Definition 4.6). It follows that

$$\Phi_t(g_0, w_0, \nu_0) = \gamma(t)(g_0, w(t), \nu(t)) = \gamma(t)(g_0, \phi_t(w_0, \nu_0))$$

where $\gamma(t) = g(t)g_0^{-1} \in g_0Z_{\mu_e, \nu_0}g_0^{-1}$. Hence $\Phi_t(W \times U_S) \subseteq A_{W,V}(\mu_e)U$, as required. $\blacksquare$

5.2 Split Momenta

Theorem 5.2 can be improved when μ_e is split. In this case Proposition 4.10 says $Z_{\mu_e, V} = G_{\mu_e}^o$ and so $A_{V, W}(\mu_e) = G_{\mu_e}^{o, W} \equiv \bigcup_{g \in W} g G_{\mu_e}^o g^{-1}$. Let $G_{\mu_e}^0 = L_{\mu_e} K_{\mu_e}$ where L_{μ_e} is a submanifold of $G_{\mu_e}^o$ and K_{μ_e} a subgroup of $G_{\mu_e}^0$ for which there exists a K_{μ_e} -invariant inner product on $\mathfrak{g}^*$. Such a splitting exists for any connected Lie group with K_{μ_e} a maximal compact subgroup of $G_{\mu_e}^o$ [13, Theorem 3.1]. However in some cases it may be possible to take K_{μ_e} larger than this. For any neighbourhood W of 1 in G define $L_{\mu_e}^W = \bigcup_{g \in W} g L_{\mu_e} g^{-1}$.

Corollary 5.3 *If p_e is a G -stable relative equilibrium and $\mu_e = J(p_e)$ is split then p_e is an $L_{\mu_e}^W K_{\mu_e}$ -stable equilibrium for every neighbourhood W of 1 in G .*

Proof. Let $\bar{K}_{\mu_e}$ denote the quotient of K_{μ_e} which acts effectively on $\mathfrak{g}^*$. The kernel T of the homomorphism $K_{\mu_e} \rightarrow \bar{K}_{\mu_e}$ acts trivially on $\mathfrak{g}^*$, and hence also on $\mathfrak{g}$. It follows that T must lie in the centre of G^o .

For any neighbourhood W of 1 in G define $K_{\mu_e}^W = \bigcup_{g \in W} g K_{\mu_e} g^{-1}$. For any neighbourhood U of p_e in P we claim that there exist neighbourhoods W of 1 in G and U' of p_e in P such that $K_{\mu_e}^W U' \subseteq K_{\mu_e} U$. Indeed consider the continuous map $\tau: G^o \times \bar{K}_{\mu_e} \times P \rightarrow P$ defined by $\tau(g, kT, p) = k^{-1} g k g^{-1} p$. Note that this is well defined because T lies in the centre of G^o . So given U and kT there exist open neighbourhoods W_k of 1 in G^o , V_k of kT in $\bar{K}_{\mu_e}$ and U_k of p in P such that $\tau(W_k, V_k, U_k) \subseteq U$. Since $\bar{K}_{\mu_e}$ is compact there exists a finite subcover $\{V_{k_i}\}_{i=1}^a$ of $\bar{K}_{\mu_e}$. Put $W = \bigcap_{i=1}^a W_{k_i}$ and $U' = \bigcap_{i=1}^a U_{k_i}$ to prove the claim.

For such a choice of U' and W we have

$$G_{\mu_e}^{o, W} U' \subseteq L_{\mu_e}^W K_{\mu_e}^W U' \subseteq L_{\mu_e}^W K_{\mu_e} U.$$

It follows that if p_e is $G_{\mu_e}^{o, W}$ -stable then it is also $L_{\mu_e}^W K_{\mu_e}$ -stable. ■

The following corollary generalises the results of [35, 19, 33] by requiring only that p_e is G -stable and not that the restriction of $d^2 H_{\xi_e}(p_e)$ to the symplectic normal space N_1 is definite.

Corollary 5.4 *If p_e is a G -stable relative equilibrium and there exists a $G_{\mu_e}^o$ -invariant inner product on $\mathfrak{g}^*$ then p_e is $G_{\mu_e}^o$ -stable.*

We end this section by applying Corollary 5.3 to the case of a semidirect product $G = K \ltimes \mathbb{V}$ with K a compact Lie group acting linearly on a finite dimensional real vector space $\mathbb{V}$. Assume that μ_e is split and G_{μ_e} has the form $G_{\mu_e} = K_{\mu_e} \ltimes \mathbb{V}_{\mu_e}$ where K_{μ_e} is a closed subgroup of K and $\mathbb{V}_{\mu_e}$ is a K_{μ_e} -invariant subspace of $\mathbb{V}$. Then $G_{\mu_e} = \mathbb{V}_{\mu_e} K_{\mu_e}$. Let W denote an open neighbourhood of $(1, 0)$ in $K \ltimes \mathbb{V}$ of the form $W_K \times W_{\mathbb{V}}$ where W_K is a neighbourhood of 1 in K and $W_{\mathbb{V}}$ is a neighbourhood of 0 in $\mathbb{V}$. Then a calculation shows $\mathbb{V}_{\mu_e}^W = W_K^{-1} \mathbb{V}_{\mu_e} \subseteq \mathbb{V}$, a generalised ‘cone’ in $\mathbb{V}$. Set theoretically $A = W_K^{-1} \mathbb{V}_{\mu_e} K_{\mu_e}$ is equal to $K_{\mu_e} \times W_K^{-1} \mathbb{V}_{\mu_e} \subseteq K \ltimes \mathbb{V}$, so we obtain the following result.

Corollary 5.5 *Let p_e be a $K \ltimes \mathbb{V}$ -stable relative equilibrium for which μ_e is split and $G_{\mu_e} = K_{\mu_e} \ltimes \mathbb{V}_{\mu_e}$. Then p_e is a $K_{\mu_e} \times W_K^{-1} \mathbb{V}_{\mu_e}$ -stable relative equilibrium for any open neighbourhood W_K of 1 in K .*

5.3 Euclidean Invariant Hamiltonian Systems

Finally, we apply the A -stability results of this section to relative equilibria of Euclidean invariant Hamiltonian systems and briefly discuss their implications for the stability of rigid bodies in fluids [17, 18]. As in Corollaries 4.18 and 4.19 we treat the cases where the symmetry group is the special Euclidean group $G = SE(2)$ of the plane or the special Euclidean group $G = SE(3)$ of three-space. We assume that the relative equilibria are G -stable and compute subsets $A \subset G$ for which the relative equilibria are A -stable.

First consider the symmetry group $G = SE(2)$. For a regular momentum value μ_e (case 1 of Corollary 4.18) let C be any open cone containing μ_e^a and $A = \{1\} \times C$. Then, for some open neighbourhood $W_{SO(2)}$ of $1 \in SO(2)$, $C \subseteq \bigcup_{R \in W_{SO(2)}} R\mathbb{R}\mu_e^a$ and so p_e is A -stable by Corollary 5.5. The cone can be made arbitrarily ‘thin’ but smaller cones will require initial conditions closer to p_e . For a non-regular momentum value (case 2 of Corollary 4.18) we have $G_{\mu_e} = G$ so trivially $A = G$ since A must contain G_{μ_e} .

Now let $G = SE(3)$. As in the case of $SE(2)$ symmetry, for a regular momentum value μ_e (case 1 of Corollary 4.19) we can apply Corollary 5.5 and conclude that p_e is A -stable for $A = SO(2) \times C$ where C is any open cone containing μ_e^a and $SO(2)$ consists of rotations about μ_e^a . For zero momentum $A = SE(3)$ trivially. Case 2 of Corollary 4.19 with $\mu_e^r \neq 0$ is more interesting because μ_e is not split and so Corollary 5.5 does not apply and a direct analysis of the sets $A_{V,W}(\mu_e) = \bigcup_{g \in W} gZ_{\mu_e,V}g^{-1}$ used in Theorem 5.2 is required. The coadjoint orbit through $(\mu_e^r, 0)$ is two sphere $\{(\mu^r, 0) : |\mu^r| = |\mu_e^r|\}$ and so a transverse section to the orbit can be taken as

$$\Sigma = \{ \mu_e + (\nu^r, \nu^a) : \nu^r = t\mu_e^r, t \in \mathbb{R}, \nu^a \in \mathbb{R}^3 \} \cong \mathbb{R} \times \mathbb{R}^3.$$

Let θ_{R,μ_e^r} denote the angle between $R\mu_e^r$ and μ_e^r . We claim that, if $\epsilon_0 > 0$ and $\epsilon_1 > 0$, and

$$A_{\epsilon_0, \epsilon_1} \equiv \{ (R, a) \in SE(3) : |\sin \theta_{R,\mu_e^r}| < \epsilon_1 |a| + \epsilon_0 \},$$

then there are neighbourhoods $V \subseteq \Sigma$ of μ_e and W of $1 \in SE(3)$ such that $A_{V,W} \subseteq A_{\epsilon_0, \epsilon_1}$. Therefore by Theorem 5.2 an $SE(3)$ stable relative equilibrium with a momentum as in Case 2 of Corollary 4.19 with $\mu_e^r \neq 0$ is $A_{\epsilon_0, \epsilon_1}$ -stable for any $\epsilon_0 > 0$ and $\epsilon_1 > 0$. To prove the claim, using the $SE(3)$ coadjoint action [26] one computes that $Z_{\mu_e, \nu}$ is the set of $(R, a) \in SE(3)$ satisfying

$$\mu_e^r \times (R(\mu_e^r + \nu^r) + a \times R\nu^a) = 0. \quad (5.8)$$

In this equation we think that $\nu^r = t\mu_e^r$ where t and ν^a are small. If a is small then $\mu_e^r \times R\mu_e^r \approx 0$, so R is nearly a rotation about μ_e^r , but the equation admits solutions for $R \in SO(3)$ arbitrary as long as a is large enough. We rewrite (5.8) as

$$(1+t)\mu_e^r \times R\mu_e^r = -\mu_e^r \times (a \times R\nu^a)$$

whereupon

$$|\sin \theta_{R,\mu_e^r}| = \frac{|\mu_e^r \times R\mu_e^r|}{|\mu_e^r|^2} \leq \frac{|\nu^a|}{|\mu_e^r|} |a|. \quad (5.9)$$

The action of conjugation is

$$(R', a') \equiv (\tilde{R}, \tilde{a})(R, a)(\tilde{R}, \tilde{a})^{-1} = (\tilde{R}R\tilde{R}^{-1}, \tilde{R}a - \tilde{R}R\tilde{R}^{-1}\tilde{a} + \tilde{a}).$$

By continuity, for arbitrary ϵ_0 there is a neighbourhood W of $1 \in SE(3)$ such that if $(\tilde{R}, \tilde{a}) \in W$ then

$$|\sin \theta_{R', \mu_e^r}| = |\sin \theta_{\tilde{R}R\tilde{R}^{-1}, \mu_e^r}| < |\sin \theta_{R, \mu_e^r}| + \frac{\epsilon_0}{2}. \quad (5.10)$$

The required R uniformity of this estimate follows from compactness of $SO(3)$. Combining (5.9) and (5.10), and choosing V so that $|\nu^a| < |\mu_e^r| \epsilon_1$ gives

$$|\sin \theta_{R', \mu_e^r}| < \frac{|\nu^a|}{|\mu_e^r|} |a| + \frac{\epsilon_0}{2} < \epsilon_1 |a| + \frac{\epsilon_0}{2}.$$

Also, by shrinking W there is the uniform (in a and R) estimate

$$|a'| = |\tilde{R}a - \tilde{R}R\tilde{R}^{-1}\tilde{a} + \tilde{a}| \geq |a| - |\tilde{R}R\tilde{R}^{-1}\tilde{a} - \tilde{a}| \geq |a| - \frac{\epsilon_0}{2\epsilon_1}$$

from which follows

$$|\sin \theta_{R', \mu_e^r}| < \epsilon_1 \left(|a'| + \frac{\epsilon_0}{2\epsilon_1} \right) + \frac{\epsilon_0}{2} = \epsilon_1 |a'| + \epsilon_0,$$

as required.

This $A_{\epsilon_0, \epsilon_1}$ -stability implies orientation stability only when translation is a priori confined. Notice that for the set A given by Corollary 5.3 for split μ_e the projection of A to the compact part of G is the same as that of G_{μ_e} . Equation 5.8 shows that in the non-split case these projections can be very different.

The results described in this section were inspired by the stability analysis of Leonard and Marsden [18] of the Kirchhoff model for the dynamics of a rigid body in a fluid. For G a semidirect product of a compact group and a vector space they prove Γ -stability results for certain groups Γ between G_{μ_e} and G . For regular momentum values of $SE(2)$ -invariant systems their group Γ is $\mathbb{R}^2$, while for regular momentum values of $SE(3)$ -invariant systems it is $SO(2) \times \mathbb{R}^3$. However numerical integration of an example with $SE(3)$ symmetry suggests that the drift in the translational direction only occurs within a cone, in accord with the results of this section.

Acknowledgments

This work was partially supported by an EPSRC Visiting Fellowship (GR/L57074) and an NSERC individual research grant for GWP, an EPSRC Research Grant (GR/K99893), a Scheme Four grant from the London Mathematical Society, a European Community Marie Curie Fellowship (HPMF-CT-2000-00542) for CW, and by European Community funding for the Research Training Network ‘MASIE’ (HPRN-CT-2000-00113). The authors thank the University of Warwick Mathematics Institute for its hospitality during several visits when parts of the paper were written.

References

- [1] V.I. Arnold. Instability of dynamical systems with several degrees of freedom. *Soviet Math.*, 5:581–585, 1964.

- [2] V.I. Arnold. On an a priori estimate in the theory of hydrodynamic stability. *Am. Math. Soc. Transl.*, 19:267–269, 1969.
- [3] V.I. Arnold. *Mathematical Methods of Classical Mechanics*. Springer-Verlag, 1978.
- [4] V.I. Arnold and B.A. Khesin. *Topological Methods in Hydrodynamics*. Springer-Verlag, 1998.
- [5] J.M. Ball and J.E. Marsden. Quasiconvexity at the boundary, positivity of the second variation and elastic stability. *Arch. Rat. Mech. Anal.*, 86:251–277, 1984.
- [6] L. Bates and E. Lerman. Proper group actions and symplectic stratified spaces. *Pac. J. Math.*, 181(2):201–229, 1997.
- [7] U. Bessi. Arnold’s diffusion with two resonances. *J. Differential Equations*, 137:211–239, 1997.
- [8] E. Brieskorn. Singular elements of semisimple algebraic groups. In *Actes Congrès Intern. Math. 1970*, t2:279–284, 1970.
- [9] S.J. Chern and J.E. Marsden. A note on symmetry and stability for fluid flows. *Geophys. Astrophys. Fluid Dynamics*, 51:19–26, 1990.
- [10] R.H. Cushman and R.M. Roberts. Poisson structures transverse to coadjoint orbits. *Bull. Sci. Math.*, to appear.
- [11] M. Duflo and M. Vergne. Une propriété de la représentation coadjointe d’une algèbre de Lie. *C.R. Acad. Sci. Paris*, 268:583–585, 1969.
- [12] V. Guillemin and S. Sternberg. A normal form for the moment map. In *Differential Geometric Methods in Mathematical Physics*, S.Sternberg (Ed.), D. Reidel, 1984.
- [13] G. Hochschild. *The Structure of Lie Groups*. Holden-Day, 1965.
- [14] D.D. Holm, J.E. Marsden, T.S. Ratiu, and A. Weinstein. Nonlinear stability of fluid and plasma equilibria. *Phys. Rev.*, 123:1–116, 1985.
- [15] P.S. Krishnaprasad. Eulerian many body problems. In J.E. Marsden, P.S. Krishnaprasad, and J.C. Simo (Ed), *Dynamics and Control of Multibody Systems*, volume 97 of *Contemporary Math.* AMS, 1989.
- [16] F. Laurent-Polz. Point vortices on the sphere: a case with opposite vorticities. *Nonlinearity*, to appear.
- [17] N.E. Leonard. Stability of a bottom-heavy underwater vehicle. *Automatica–J. IFAC*, 33:331–346, 1997.
- [18] N.E. Leonard and J.E. Marsden. Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry. *Physica D*, 105:130–162, 1997.
- [19] E. Lerman and S.F. Singer. Stability and persistence of relative equilibria at singular values of the moment map. *Nonlinearity*, 11:1637 – 1649, 1998.
- [20] D. Lewis, T.S. Ratiu, J.C. Simo and J.E. Marsden. The heavy top: a geometric treatment. *Nonlinearity*, 5:1–48, 1992.
- [21] D. Lewis and J.C. Simo. Nonlinear stability of rotating pseudo-rigid bodies. *Proc. Roy. Soc. London Ser. A*, 427:281–319, 1990.
- [22] P. Libermann and C.-M. Marle. *Symplectic Geometry and Analytical Mechanics*. Reidel, 1987.
- [23] S. Lie. Theorie der Transformationsgruppen. Zweiter Abschnitt. Teubner, 1890.
- [24] C. Marchioro and M. Pulvirenti. *Mathematical Theory of Incompressible Nonviscous Fluids*. Springer-Verlag, 1994.
- [25] C.-M. Marle. Modèle d’action hamiltonienne d’un groupe de Lie sur une variété symplectique. *Rendiconti del Seminario Matematico, Università e Politecnico, Torino*, 43:227–251, 1985.
- [26] J.E. Marsden and T.S. Ratiu. *Introduction to Mechanics and Symmetry*, volume 17 of *Texts in Applied Math.* Springer-Verlag, 1994.
- [27] J.E. Marsden, J.C. Simo, D. Lewis, and T.A. Posbergh. Block diagonalization and the energy-momentum method. In J.E. Marsden, P.S. Krishnaprasad, and J.C. Simo (Ed.), *Dynamics and Control of Multibody Systems*, Volume 97 of *Contemporary Math.*, 297–313. AMS, 1989.
- [28] J.E. Marsden and A. Weinstein. Reduction of symplectic manifolds with symmetry. *Rep. Math. Phys.*, 5:121–130, 1974.

- [29] E.T. Matsui. An equivariant Liapunov stability test and the energy-momentum-Casimir method. *Preprint*, 2001.
- [30] A. Mielke. *Hamiltonian and Lagrangian Flows on a Center Manifold with Applications to Elliptic Variational Problems*, Volume 1489 of *Lect. Notes Math.* Springer-Verlag, 1991.
- [31] J. Montaldi. Persistence and stability of relative equilibria. *Nonlinearity*, 10:449–466, 1997.
- [32] P. Newton. *The N-Vortex Problem*, Volume 145 of *Appl. Math. Sc.* Springer-Verlag, 2001.
- [33] J.-P. Ortega and T.S. Ratiu. Stability of Hamiltonian relative equilibria. *Nonlinearity*, 12:693–720, 1999.
- [34] G.W. Patrick. The dynamics of two coupled rigid bodies in three space. In J.E. Marsden, P.S. Krishnaprasad, and J.C. Simo (Eds.), *Dynamics and Control of Multibody Systems*, Volume 97 of *Contemporary Math.*, 297–313. AMS, 1989.
- [35] G.W. Patrick. Relative equilibria in Hamiltonian systems: the dynamic interpretation of nonlinear stability on a reduced phase space. *J. Geom. Phys.*, 9:111–119, 1992.
- [36] G.W. Patrick. Dynamics of perturbed relative equilibria of point vortices on the sphere or plane. *J. Nonlin. Sci.*, 10:401–415, 2000.
- [37] G.W. Patrick, R.M. Roberts and C. Wulff. Linear and nonlinear stability of Hamiltonian relative equilibria. *In preparation*, 2001.
- [38] G.W. Patrick, R.M. Roberts and C. Wulff. Stability of nongeneric relative equilibria of underwater vehicles. *In preparation*, 2001.
- [39] S. Pekarsky and J.E. Marsden. Point vortices on a sphere: stability of relative equilibria. *J. Math. Phys.*, 39:5894–5906, 1998.
- [40] T.A. Posbergh, J.C. Simo, and J.E. Marsden. Stability analysis of a rigid body with attached geometrically nonlinear rod by the energy momentum method. In J. E. Marsden, P. S. Krishnaprasad, and J. C. Simo (Eds.), *Dynamics and Control of Multibody Systems*, Volume 97 of *Contemporary Math.*, 371–397. AMS, 1989.
- [41] R.M. Roberts, C. Wulff and J.S.W. Lamb. Hamiltonian systems near relative equilibria. *J. Differential Equations*, to appear.
- [42] J.C. Simo, D.R. Lewis and J.E. Marsden. Stability of relative equilibria I: The reduced energy momentum method, *Arch. Rat. Mech. Anal.*, 115:15–59, 1991.
- [43] J.C. Simo, T. Posbergh, and J.E. Marsden. Stability of coupled rigid body and geometrically exact rods: Block diagonalization and the energy-momentum method. *Phys. Rep.*, 193:279–360, 1990.
- [44] J.C. Simo, T.A. Posbergh and J.E. Marsden. Stability of relative equilibria II: Three dimensional elasticity, *Arch. Rat. Mech. Anal.*, 115:61–100, 1991.
- [45] P. Slodowy. *Simple Singularities and Simple Algebraic Groups*, Volume 815 of *Lect. Notes Math.* Springer-Verlag, 1980.
- [46] Y.-H. Wan. The stability of rotating vortex patches. *Commun. Math. Phys.*, 107:1–20, 1986.
- [47] A. Weinstein. Local structure of Poisson manifolds. *J. Differential Geom.*, 18:523–557, 1983. Errata and addenda. *J. Differential Geom.*, 22:255, 1985.
- [48] A. Weinstein. Stability of Poisson-Hamilton equilibria. In J.E. Marsden (Ed.), *Fluids and Plasmas: Geometry and Dynamics*, Volume 28 of *Contemporary Math.*, 3–13. AMS, 1984.
- [49] C. Wulff. Persistence of relative equilibria in Hamiltonian systems with non-compact symmetry. *Preprint*, 2001.
- [50] C. Wulff, G.W. Patrick and R.M. Roberts. Stability of Hamiltonian Relative Equilibria with Energy Methods. In D. Bambusi, G. Gaeta and M. Cadoni (Eds.), *Proceedings of “Symmetry and Perturbation Theory 2001”*. World Scientific, 2001.